TITLE: Structure and antioxidant activity of Maillard reaction products from α -lactalbumin and β -lactoglobulin with ribose in an aqueous model system

AUTHORS: Zhanmei Jiang, André Brodkorb

This article is provided by the author(s) and Teagasc T-Stór in accordance with publisher policies.

Please cite the published version.

The correct citation is available in the T-Stór record for this article.

NOTICE: This is the author's version of a work that was accepted for publication in *Food Chemistry*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Food Chemistry*, Volume 133, Issue 3, 1 August 2012, Pages 960–968. DOI: 10.1016/j.foodchem.2012.02.016

This item is made available to you under the Creative Commons Attribution-Non commercial-No Derivatives 3.0 License.

Structure and antioxidant activity of Maillard reaction products from α -lactalbumin and β -lactoglobulin with ribose in aqueous model system

Zhanmei Jiang^a, André Brodkorb^{b,*}

^a Key Laboratory of Dairy Science, Ministry of Education, Northeast Agricultural University, Harbin 150030, China; ^b Teagasc Food Research Centre, Moorepark, Fermoy, Co. Cork, Ireland

*Corresponding Author: Dr. André Brodkorb, Teagasc Food Research Centre, Moorepark, Fermoy, Co. Cork, Ireland.

E-mail: andre.brodkorb@teagasc.ie, Tel: +353-(0)25-42222, Fax: +353-(0)25-42340

Abstract

Maillard reaction products (MRPs) were prepared from aqueous model mixtures containing 3% (w/w) ribose and 3% (w/w) α -lactalbumin (α -LA) or β -lactoglobulin (β -LG), heated at 95°C, for up to 5 hours. The pH of MRPs decreased significantly during heat treatment of α -LA-Ribose and β -LG-Ribose mixtures from 8.4 to 5.3. The amino group content in MRPs derived from the α -LA-Ribose and β -LG-Ribose model system was decreased noticeably during the first hour and did not change thereafter. The loss of free ribose in MRPs was higher for β -LG-Ribose than for α -LA-Ribose. During the Maillard Reaction, the concentration of native and non-native α -LA or β -LG decreased and the formation of aggregates was observed. Fluorescence intensity of the β -LG-Ribose MRPs reached the maximum within 1 hour, compared to 2 hours for α -LA-Ribose MRPs. Meanwhile, modification of the UV/vis absorption spectra for α -LA and β -LG was mainly due to a condensation reaction with ribose. Dynamic light scattering showed a significant increase in the particle size of the MRPs. Size exclusion chromatography of MRPs revealed the production of both high and low molecular weight material. Electrophoresis of MRPs indicated polymerization of α -LA and β -LG monomers via inter-molecular disulfide bridge but also via other covalent bonds. MRPs from α -LA-Ribose and β -LG-Ribose exhibited increased antioxidant activities, therefore these MRPs may be used as natural antioxidants in food products.

Key words: Maillard reaction, α -Lactalbumin, β -Lactoglobulin, Ribose, Antioxidant Activity

46 **Introduction**

47 Maillard reaction is one of the most important chemical reactions during food preparation (James P,
48 1986; Jennifer M, 1990; Werner, 1982). Maillard reaction involves a series of reactions steps involving
49 the condensation between carbonyl groups and amino groups, resulting in a large number of the
50 Maillard reaction products (MRPs), such as volatile compounds, non-volatile intermediates and dark-
51 brown melanoidins (Ledl, Beck, Sengl, Osiander, Estendorfer, Severin, et al., 1989; Van Boekel, 1998).
52 Although Maillard reaction is of a great importance in production of aroma, taste and colour, it can
53 result a loss of nutritional value, digestive incompatibility and potential toxicity. Characterization of the
54 MRPs could help controlling the formation of advanced Maillard reaction products (noxious effects in
55 diabetes and in age-related cardiovascular diseases). However, controlled reactions can also modify
56 protein functionality (Hiller & Lorenzen, 2010; Katayama & Saeki, 2004; Li, Lu, Luo, Chen, Mao,
57 Shoemaker, et al., 2009; Oliver, Melton, & Stanley, 2006; Qi, Yang, & Liao, 2009; Wahyuni, Ishizaki,
58 & Tanaka, 1998; Wooster & Augustin, 2006).

59 Whey proteins are extensively utilized in the food processing. α -lactalbumin (α -LA) and β -
60 lactoglobulin (β -LG) are two protein major components in whey and play important roles in the
61 functional properties and physiological function of whey proteins (Chatterton, Smithers, Roupas, &
62 Brodkorb, 2006; Croguennec, Renault, Bouhallab, & Pezennec, 2006; Rojas, Goff, Senaratne,
63 Dalglish, & Flores, 1997). During recent years, increasing attention has been directed towards the
64 utilization of Maillard reaction to improve functional properties of α -LA or β -LG including
65 emulsifying properties, foaming properties and heat stability (Aoki, Kitahata, Fukumoto, Sugimoto,
66 Ibrahim, Kimura, et al., 1997; Chevalier, Chobert, Popineau, Nicolas, & Haertlé, 2001; Medrano,
67 Abirached, Panizzolo, Moyna, & Añón, 2009; Wooster & Augustin, 2006). Some reports focused on
68 mechanism and chemical properties of Maillard reaction from α -LA or β -LG and sugars. For example,
69 characteristics of conjugates between β -LG and acacia gum (Schmitt, Bovay, & Frossard, 2005) or
70 chitosan (Miralles, Martínez-Rodríguez, Santiago, van de Lagemaat, & Heras, 2007) were investigated
71 upon for dry -state model systems. Some structural characterization of MRPs of glycated β -LG with
72 galactose and tagatose has been carried out under dry heat condition (Corzo-Martinez, Moreno, Olano,
73 & Villamiel, 2008). Above-mentioned studies on MRPs of α -LA or β -LG were only based on dry
74 heating and relatively mild reaction condition. However, during the food processing, Maillard reaction
75 generally happens in hydrated or aqueous system under high temperature conditions. However, only

few published examples are available to date. Meanwhile, few studies were carried out on the antioxidant properties of Maillard reaction products of α -LA or β -LG. Conjugates of α -LA with three monosaccharides produced in the dry state showed antioxidant activities (Sun, Hayakawa, Puangmanee, & Izumori, 2006). However, the MRPs were not fully characterised, the objectives of this study were to describe the structural changes and antioxidant activity during Maillard reaction of α -LA and β -LG with ribose in an aqueous system. Therefore conditions were chosen to promote Maillard reaction i.e. high heat and pH as well as the choice of ribose as the one of the most reactive monosaccharides.

2. Materials and methods

2.1. Materials

Commercial α -lactalbumin (95% protein content, approximately 3.0% β -lactoglobulin) and β -lactoglobulin (94% protein content, approximately 3.5% α -lactalbumin) were purchased from Davisco Foods International, Inc. (Eden Prairie, MN, USA). D-Ribose, OPA (o-phthalaldehyde), L-leucine, Potassium ferricyanide(III), 2,2-diphenyl-1-picryl-hydrazyl (DPPH), sodium dodecyl sulphate (SDS), β -mercaptoethanol, L-leucine, and trifluoroacetic acid (TFA) and other reagents were purchased from Sigma Aldrich Co. (St. Louis, MO, USA) unless stated otherwise. All chemicals used in this study were of analytical grade.

2.2. Preparation of α -LA-Ribose and β -LG-Ribose MRPs

α -LA and β -LG were dissolved in Milli-Q[®] water at a protein concentration of 30 g/L with the ribose (30 g/L), respectively. The initial pH of the mixtures was adjusted to 8.4 using pH meter (pH 340i, WTW, 82362, Germany). The mixtures were then transferred to sealed screw-top test tubes, and heated in water bath at 95°C. Samples were taken after heating for 0, 0.5, 1, 2, 3, 4, and 5 hours and immediately submerged in ice water. Control experiments were also carried out with no added ribose.

2.3. Determination of free amino groups

The quantity of free amino groups was determined by the OPA method (Church, Swaisgood, Porter, & Catignani, 1983). MRPs (35 μ L, 12-fold dilution) were mixed with 1 mL of OPA reagent. After vortexing and a minimal 5 min delay in the dark at room temperature, the absorbance was recorded at 340 nm. Absorbance readings were converted into free amino contents using a calibration curve obtained with L-leucine (0 to 0.5 g/L) as a standard. The blank was determined in the same manner, except that MilliQ[®] water was used instead of sample. Data were finally expressed as relative concentrations (%) in comparison with the content of unheated samples.

2.4. Determination of ribose in MRPs

The free ribose in MRPs samples were determined by HPLC (waters 2695 Alliance, Waters Inc., USA) equipped with a WATO44355 carbohydrate column (4.6×250 mm I.D., 4 µm particle size, Waters, Ireland). 50µL of sample were injected and eluted at a flow rate of 0.5mL/min using a mobile phase of 9mM H₂SO₄. The column temperature was set at 60°C. A refractive index detector (2414 Waters Inc., USA) was used for sample detection. Data analysis and quantification were performed using Empower[®] Chemstation software. Changes in ribose were expressed as relative concentrations (%) in comparison with the original ribose content in non-heated samples.

2.5. Size exclusion chromatography (SEC)

The MRP samples were separated by size exclusion chromatography using a series connection column with TSK G 3000SW XL (7.8mm ID×30cm L, 5µm) and TSK G 2000SW XL (7.8mm ID×30cm L, 5µm, TosoHaas, Montgomeryville, PA, USA) fitted to a TSK guard column (7.5mm×7.5cm). Samples (20 µL of 2.5 g/L protein) were eluted with 50 mM Tris-HCl buffer (pH 7.0) was used as the eluent at a flow rate of 0.6 mL/min. The absorbance was monitored at 280nm. 5mL of the MRP samples were added 5mL of 0.01M acetate buffer pH 4.6. The mixtures were centrifuged for 20 min at 5,000g in order to get the supernatant. The native protein concentration in MRPs was determined by adding 0.01M acetate buffer pH 4.6 analysis of the supernatant by SEC. The method was calibrated using α-LA and β-LG standards.

2.6. Determination of Native, Non-native and Aggregated α-LA and β-LG concentrations in MRPs

The concentration of native protein was determined as described above. Monomeric protein (including both native and non-native) were directly obtained using SEC. The concentrations of non-native monomeric α-LA and β-LG were calculated by subtraction of the level of native protein from that of monomeric protein. The amount of aggregation was calculated by subtraction of the amount of monomer (SEC) from the initial protein concentration.

2.7. Absorption and fluorescence spectroscopy

UV/vis absorption spectra were recorded using a CARY 1 UV-visible spectrophotometer (Varian, USA) with the wavelength ranging from 200 to 800 nm. The α-LA-Ribose and β-LG-Ribose model MRPs were diluted (1:120, v/v) with Milli-Q[®] water for UV/vis analysis.

Fluorescence spectra of α-LA-Ribose and β-LG-Ribose MRPs were determined using a temperature controlled Cary Eclipse Fluorescence Spectrometer (Varian, USA). Samples were diluted (1:120 v/v)

with MilliQ[®] water. For fluorescence excitation spectra, the emission wavelength was to 420nm and the excitation was scanned from 300 to 400 nm. For fluorescence spectra, the excitation wavelength was 337 nm, and the emission was scanned from 370 to 570 nm. Both excitation and emission slits were set to 10 nm, the temperature was 20 °C, and the scan rate was 120 nm/min.

2.8. Dynamic Light Scattering (DLS)

The hydrodynamic diameter of α -LA-Ribose and β -LG-Ribose MRPs were determined using a Zetasizer Nano system (Malvern Instruments Inc., Worcester, UK). The measurements were carried out at 25 °C using a scattering angle of 12° and the viscosity of the solvent was assumed to be the same as water.

2.9. Sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE)

SDS-PAGE was performed according to a modified method of Laemmli (1970), using 4% acrylamide stacking gel and 15% acrylamide separating gel with a vertical gel electrophoresis cell (Bio-Rad, Richmond, CA, USA). MRP samples were prepared under both reducing (with β -mercaptoethanol) and non-reducing conditions. Gels were stained with 0.5% Coomassie Brilliant Blue R-250. Destaining was conducted with a solution of isopropanol and acetic acid. Molecular weight markers (10 to 250 kDa) (Bio-Rad Laboratories, CA, USA) were used for M_w estimation of samples.

2.10. Determination of reducing power

The reducing power of samples of α -LA-Ribose and β -LG-Ribose MRPs was determined according to the method of Chawla, Chander and Sharma (2009) with some modifications. 0.1mL of MRPs sample (12-fold dilution) was mixed with 0.5 mL of 0.2 M sodium phosphate buffer (pH 6.6) and 0.5 mL of 1% potassium ferricyanide. The reaction mixtures were incubated in a water bath at 50°C for 20 min, followed by addition of 0.5 mL of 10% trichloroacetic acid. The mixtures were then centrifuged at 5,000g using a Eppendorf 5417R centrifuge (Eppendorf AG, Hamburg, Germany) for 10 min at 25 °C. The supernatant (0.5 mL) was treated with 0.5 mL of Milli-Q[®] water and 0.1mL of 0.1% FeCl₃. After the reaction, the mixture was left for 10 min, the absorbance was measured at 700 nm. Higher absorbance of the reaction mixture indicated greater reducing power.

2.11. Radical scavenging activity using the 2,2-Diphenyl-1-picryl-hydrazyl (DPPH)

DPPH radical scavenging activity of the MRPs was estimated according to a modified method of Yen and Hsieh (1995). An aliquot of MRP sample (250 μ L) was added to 1 mL of a solution of DPPH, prepared freshly, at a concentration of 0.1mM in ethanol. The reaction solution was then mixed

vigorously and allowed to stand at room temperature for 30 min in the dark. The mixture was centrifuged for 10 min at 4,000g. The absorbance of the supernatant was measured at 517 nm using a CARY 1 UV-visible spectrophotometer. The percentage of DPPH radical scavenging activity was calculated as follows:

$$\text{DPPH radical scavenging activity (\%)} = (1 - \frac{A_i - A_j}{A_c}) \times 100\%$$

Where A_i is the absorbance of sample, A_j is the absorbance of ethanol added instead of 0.1 mM DPPH and A_c is the absorbance of ethanol added instead of MRP samples.

2.12 Statistical analysis

Analyses including the determination of pH, free amino group, ribose, reducing power and DPPH radical scavenging activity were run in at least triplicate. Results were expressed as means with the standard deviation. Analysis of variance (ANOVA) was performed and significant differences among means from triplicate or more by the Duncan's multiple range tests using the SPSS system software 13.0. For the DLS analysis, the experiments were carried out in duplicates and each measurement was made in triplicate. SEC measurements were carried out in duplicates.

3. Results and Discussion

3.1. Changes in pH

The changes in the pH of MRPs derived from aqueous α -LA-Ribose and β -LG-Ribose systems as a function of heating time were shown in Fig.1A. The pH of α -LA-Ribose and β -LG-Ribose MRPs decreased significantly as the heating time increased up to 5 h ($P < 0.05$). During the first 2 hours, the pH decreased at a higher rate, and the pH of β -LG-Ribose MRPs decreased more quickly than α -LA-Ribose MRPs ($P < 0.05$). After 2 hours, the pH of α -LA-Ribose and β -LG-Ribose MRPs slightly decreased up to 5 hours and there was no obvious difference in the change of pH between two systems. The result was in accordance with Benjakul, Lertittikul and Bauer (2005) and Gu, Kim, Hayat, Xia, Feng and Zhang (2009) who found that the pH of protein-sugar MRPs decreased with the heating time and levelled off with prolonged heating. The reduction in pH is generally attributed to the formation of organic acids, such as acetic acid and formic acid in the intermediate MRPs (Brands & van Boekel, 2002; Chen, Jin, & Chen, 2005; Rufian-Henares, Delgado-Andrade, & Morales, 2006). In addition, the consumption of amino groups and the formation of acidic compounds could also contribute to the decrease in pH. In contrast to this, the pH in α -LA and β -LG heated alone (95°C), only slightly

decreased during heating, possibly due to the thermal protein degradation (Lan, Liu, Xia, Jia, Mukunzi, Zhang, et al., 2010). Therefore, it was assumed that formation of organic acids from Maillard reaction was a key factor in the pH decrease of MRPs.

3.2. Changes in the protein amino groups

During Maillard reaction, available amino groups of the proteins contain terminal α -amino groups of protein and ϵ -amino groups of lysine residues. To assay amino group availability, the quantities of amino groups in α -LA and β -LG heated in the presence and absence (control sample) of ribose for up to 5 h were measured using the OPA method and results were shown in the Fig.1B. When α -LA and β -LG was heated without ribose at pH 8.4, the free amino groups did not significantly change during the whole heating time ($P>0.05$). Similar results were reported previously (Sun, Hayakawa, Puangmanee, & Izumori, 2006), where free amino groups of heated control α -LA did not significantly change during the 48h incubation. However, compared to the initial α -LA-Ribose and β -LG-Ribose MRPs, a significant decrease ($P<0.05$) in free amino group content, $32.52\pm0.81\%$ and $28.32\pm1.11\%$ respectively, was observed in the α -LA-Ribose and β -LG-Ribose MRPs when heated for one hour. Then the amino group loss did not changed significantly with the prolonged heating time to 5h. This indicates that heating quickly induced interaction of free amino groups, such as α -NH₂ and ϵ -NH₂ of lysine residues, with ribose by Maillard reaction within 1 hour. The pH decrease of protein-sugar systems and decrease in reactivity of the protein resulted in only a small change in the free amino group consumption during prolonged heating.

3.3. Changes of ribose and protein in MRPs

The loss of ribose and protein in the α -LA-Ribose and β -LG-Ribose MRPs, were given in Fig.1C and D as a function of the heating time. A sharp decrease in ribose concentration in both MRP model systems was observed for up to 5 h of the heating time ($P<0.05$). Within the first 3 hour, there was no statistical significance in the loss of ribose between the α -LA-Ribose and the β -LG-Ribose ($P>0.05$). However, from 4 h to 5h of heating, the loss of ribose was higher in β -LG-Ribose than in α -LA-Ribose. Sugar consumption is a critical property to evaluate the reactivity of sugars in the Maillard reaction. It was also noted that during the Maillard reaction, ribose was the most active among five tested sugars (Xylose, Arabinose, Glucose and Fructose) (Laroque, Inisan, Berger, Vouland, Dufossé, & Guérard, 2008). In the present study, ribose consumption from the α -LA-Ribose and the β -LG-Ribose was $35.74\pm1.51\%$ and $39.26\pm0.25\%$ at 5 h, respectively, compared to the initial α -LA-Ribose and the β -

LG-Ribose. 12 hours of heating led to 30% of remaining glucose when reacted with different amino acids (Kwak & Lim, 2004). Kim and Lee (2009) also reported that the loss of glucose in MRPs derived from the Glucose-Glycine, Glucose-Diglycine and Glucose-Triglycine model system was a function of heating time. The distribution of native, non-native and aggregated proteins in α -LA-Ribose and β -LG-Ribose reaction mixtures is shown in the Fig.1C and Fig.1D. Prior to heating, α -LA-Ribose MRPs model system contained 73.73% native protein and 26.27% non-native protein, but β -LG-Ribose MRPs contained 55.98 % native protein, 15.36% non-native protein and 28.66% aggregates, which is within acceptable limits for commercial α -LA and β -LG, probably due to slow denaturation and aggregation in the powders. Native protein concentration in the α -LA-Ribose and β -LG-Ribose MRPs decreased gradually and whereas aggregate increased quickly with increase heating time up to 5 h. After 0.5h of heat-treatment, the concentrations of non-native protein, which probably included those previously described (Croguennec, Bouhallab, Molle, O'Kennedy, & Mehra, 2003; Kehoe, Wang, Morris, & Brodkorb, 2011), were highest and decreased thereafter. The above results suggested that Maillard reaction of α -LA and β -LG with ribose resulted in substantial amounts of aggregates.

3.4. Spectroscopic analysis

UV/vis absorption, fluorescence excitation and emission spectra of α -LA-Ribose and β -LG-Ribose model system are shown in the Fig.2. The appearance of new compounds with an excitation maximum of $337\pm 1\text{nm}$ and an emission maximum of $420\pm 1\text{nm}$, was observed within the first 1 to 2 hours (faster for β -LG-Ribose than α -LA-Ribose) and followed by a rapid decline. The corresponding fluorescence reached a maximum value within 2 hours for α -LA-Ribose MRPs, compared to 1 hours for β -LG-Ribose MRPs. The fluorescence is associated with the early stage of the Maillard reaction and the development of fluorescent compounds, which are considered to be precursors of brown pigments formed during Maillard reaction (Morales, Romero, & JimenezPerez, 1996). The development of fluorescent compounds was found to increase with longer heating times of the rice protein and glucose (Li, Lu, Luo, Chen, Mao, Shoemaker, et al., 2009). In contrast, fluorescence was not found to increase linearly with heating time in ribose-casein systems (Jing & Kitts, 2004).

The changes in the UV/vis absorption spectra of α -LA-Ribose and β -LG-Ribose MRPs as a function of the heating time are shown in Fig. 2A' and 2B'. The absorbance of MRP samples increased significantly as the heating time increased for up to 5 h. This results are in accordance with previous observations (Kim & Lee, 2009).

3.5. Dynamic Light Scattering (DLS)

Particle size of α -LA-Ribose and β -LG-Ribose (Fig.3.) increased as a function of heating time, β -LG-Ribose Particles being larger than those of α -LA-Ribose. However, under the experimental condition (pH 8.4), particle sizes of heated α -LA and β -LG exhibited only marginal changes.

3.6. SDS-PAGE

All of α -LA-Ribose and β -LG-Ribose MRPs samples were analyzed by both non-reducing and reducing SDS-PAGE. The noncovalent interactions within aggregates are generally disrupted in the presence of SDS (non-reducing conditions) whereas addition of β -mercaptoethanol cleaves all disulphide bonds (reducing conditions).

α -LA-ribose: Both non-reducing and reducing SDS-PAGE (Fig. 4A and 4B, respectively), the amount of aggregates increased with the prolonged heat treatment. Comparing the amounts of aggregate on non-reducing and reducing SDS-PAGE, it was estimated that some aggregates were linked by disulfide bonds. However, reducing conditions failed to dissociate aggregates, suggesting other covalent, inter-molecular bonds, possibly conjugates of proteins and ribose. A gradual increase in conjugate size, from monomer to dimers and trimers etc. could be detected. Several bands between approximately 26 to 34kDa are clearly visible under non-reducing conditions, suggesting that in addition to homo-conjugates of α -LA with α -LA (M_w of 28kDa), hetero-conjugates such as α -LA with β -LG (M_w of 32kDa) could be produced. After 5 hours heat-treatment, no remaining monomeric α -LA was detected.

β -LG-ribose: Similar observations were made for β -LG-Ribose (Fig. 4 C and D). However, polymerisation appeared to be faster than for α -LA-ribose. After 4 hours heat-treatment, practically no remaining monomeric β -LG was detected. Under non-reducing condition monomers, dimers and trimers of β -LG were observed prior heating, which indicated that initial β -LG contained disulphide-linked aggregates, which was also confirmed by the above-mentioned structural characterisation (Fig.1D).

3.7. Size exclusion chromatography (SEC)

Molecular rearrangements in α -LA-Ribose and β -LG-Ribose MRPs could be observed by size exclusion chromatography (Fig.5.). It showed a decrease in the amount of monomer and an increase in aggregated α -LA and β -LG. The appearance of a new peak corresponded to small molecules. The amount of both aggregates and small molecular weight (M_w) material increased with increasing heating time; accordingly, the amount of monomer decreased. Meanwhile, the production rate and

concentrations of smaller compounds of MRPs derived from β -LG were quicker and higher than those from α -LA. α -LA and β -LG were probably covalently attached to ribose or ribose degradation products, which resulted in high M_w MRPs (melanoproteins). Meanwhile, caramelisation of ribose heated at high pH or degradation products of Maillard reaction may induce the formation of low and medium M_w compounds.

3.8. Antioxidant properties

3.8.1 DPPH radical scavenging activity

DPPH radical-scavenging activity of α -LA-Ribose and β -LG-Ribose MRPs increased significantly ($P < 0.05$) for the first two hours of heating but levelled off at 4 to 5 hours of heating, as shown in the Fig. 5A. There was no significant difference in DPPH radical-scavenging activity between α -LA-Ribose and β -LG-Ribose as the heating time increased ($P > 0.05$). In contrast to this, heated α -LA and β -LG lacked the DPPH radical-scavenging activity of its corresponding MRPs. It is recognised that DPPH radicals can be scavenged by MRPs through donation of hydrogen to form stable DPPH-H (Matthaus, 2002). The present results were in agreement with previous observations where porcine plasma MRPs of protein-reducing sugars (Benjakul, Lertittikul, & Bauer, 2005) and ultra-filtered MRPs from a casein-glucose model system showed strong DPPH radical-scavenging activity (Gu, Kim, Hayat, Xia, Feng, & Zhang, 2009).

3.8.2 Reducing power of MRPs

Reducing power of α -LA-Ribose and β -LG-Ribose MRPs increased significantly as the heating time increased ($P < 0.05$), as shown by an increase in absorbance at 700 nm (Fig. 6B). Reducing power of two model systems also increased notably within the first 2 hours but levelled off at after 5 hours of heating. Reducing power of β -LG-Ribose MRPs was significantly higher than that of α -LA-Ribose MRPs ($P < 0.05$). However, it was also shown that there was a somewhat smaller reducing power of heated α -LA and β -LG. Hydroxyl groups and pyrrole groups of MRPs may play a drinving role in reducing activity (Yanagimoto, Lee, Ochi, & Shibamoto, 2002; Yoshimura, Iijima, Watanabe, & Nakazawa, 1997). Heat-induced MRPs from casein/glucose (Gu, Abbas, & Zhang, 2009; Gu, Kim, Abbas, Zhang, Xia, & Chen, 2010) and porcine plasma protein/glucose models (Benjakul, Lertittikul, & Bauer, 2005) also exhibited reducing power.

4. Conclusion

MRPs of α -LA and β -LG with ribose were prepared under high temperature and alkali pH conditions. Free amino groups, pH and free ribose decreased considerably with increasing heat. Characteristics of Maillard reaction of α -LA-Ribose and β -LG-Ribose model system indicated production of new smaller molecules, molecular rearrangements and polymer production occurred with heat treatment increased. Based on the reducing power and radical-scavenging activity of the α -LA-Ribose and β -LG-Ribose MRPs, they may be considered as possible antioxidants for food products.

Acknowledgements

The studies were supported by Program for New Century Excellent Talents in Heilongjiang Province University (1251-NCET-009) and the Innovative Research Team of Higher Education of Heilongjiang Province (2010td11). The studies were also supported by the food program of the Teagasc Food Research Centre, Moorepark.

Reference

- Aoki, T., Kitahata, K., Fukumoto, T., Sugimoto, Y., Ibrahim, H. R., Kimura, T., Kato, Y., & Matsuda, T. (1997). Improvement of functional properties of β -lactoglobulin by conjugation with glucose-6-phosphate through the Maillard reaction. *Food Research International*, 30(6), 401-406.
- Benjakul, S., Lertittikul, W., & Bauer, F. (2005). Antioxidant activity of Maillard reaction products from a porcine plasma protein-sugar model system. *Food Chemistry*, 93(2), 189-196.
- Brands, C. M. J., & van Boekel, M. (2002). Kinetic modeling of reactions in heated monosaccharide-casein systems. *Journal of Agricultural and Food Chemistry*, 50(23), 6725-6739.
- Chatterton, D. E. W., Smithers, G., Roupas, P., & Brodkorb, A. (2006). Bioactivity of β -lactoglobulin and α -lactalbumin—Technological implications for processing. *International Dairy Journal*, 16(11), 1229-1240.
- Chen, S.-L., Jin, S.-Y., & Chen, C.-S. (2005). Relative reactivities of glucose and galactose in browning and pyruvaldehyde formation in sugar/glycine model systems. *Food Chemistry*, 92(4), 597-605.
- Chevalier, F., Chobert, J.-M., Popineau, Y., Nicolas, M. G., & Haertlé, T. (2001). Improvement of functional properties of β -lactoglobulin glycosylated through the Maillard reaction is related to the nature of the sugar. *International Dairy Journal*, 11(3), 145-152.

344 Church, F. C., Swaisgood, H. E., Porter, D. H., & Catignani, G. L. (1983). Spectrophotometric Assay
 345 Using o-Phthaldialdehyde for Determination of Proteolysis in Milk and Isolated Milk Proteins.
 346 *Journal of Dairy Science*, 66(6), 1219-1227.

347 Corzo-Martinez, M., Moreno, F. J., Olano, A., & Villamiel, M. (2008). Structural characterization of
 348 bovine beta-lactoglobulin-galactose/tagatose Maillard complexes by electrophoretic,
 349 chromatographic, and spectroscopic methods. *Journal of Agricultural and Food Chemistry*,
 350 56(11), 4244-4252.

351 Croguennec, T., Renault, A., Bouhallab, S., & Pezenec, S. (2006). Interfacial and foaming properties
 352 of sulfhydryl-modified bovine β -lactoglobulin. *Journal of Colloid and Interface Science*, 302(1),
 353 32-39.

354 Croguennec, T., Bouhallab, S., Molle, D., O'Kennedy, B. T., & Mehra, R. (2003). Stable monomeric
 355 intermediate with exposed Cys-119 is formed during heat denaturation of beta-lactoglobulin.
 356 *Biochemical and Biophysical Research Communications*, 301(2), 465-471

357 Gu, F.L., Abbas, S., & Zhang, X.M. (2009). Optimization of Maillard reaction products from casein-
 358 glucose using response surface methodology. *Lwt-Food Science and Technology*, 42(8), 1374-
 359 1379.

360 Gu, F.L., Kim, J. M., Abbas, S., Zhang, X. M., Xia, S.Q., & Chen, Z.X. (2010). Structure and
 361 antioxidant activity of high molecular weight Maillard reaction products from casein-glucose.
 362 *Food Chemistry*, 120(2), 505-511.

363 Gu, F.L., Kim, J. M., Hayat, K., Xia, S., Feng, B., & Zhang, X.M. (2009). Characteristics and
 364 antioxidant activity of ultrafiltrated Maillard reaction products from a casein-glucose model
 365 system. *Food Chemistry*, 117(1), 48-54.

366 Hiller, B., & Lorenzen, P. C. (2010). Functional properties of milk proteins as affected by Maillard
 367 reaction induced oligomerisation. *Food Research International*, 43(4), 1155-1166.

368 James P, D. (1986). Maillard Reactions: Nonenzymatic Browning in Food Systems with Special
 369 Reference to the Development of Flavor. In E. M. M. C.O. Chichester & B. S. Schweigert
 370 (Eds.), *Advances in Food Research*, vol. Volume 30 (pp. 77-138): Academic Press.

371 Jennifer M, A. (1990). Control of the Maillard reaction in food systems. *Trends in Food Science &*
 372 *Technology*, 1(0), 150-154.

373 Jing, H., & Kitts, D. D. (2004). Chemical characterization of different sugar-casein Maillard reaction
 374 products and protective effects on chemical-induced cytotoxicity of Caco-2 cells. *Food and*
 375 *Chemical Toxicology*, 42(11), 1833-1844.

376 Katayama, S., & Saeki, H. (2004). Cooperative effect of relative humidity and glucose concentration on
 377 improved solubility of shellfish muscle protein by the Maillard reaction. *Fisheries Science*,
 378 70(1), 159-166.

379 Kehoe, J., Wang, L., Morris, E., & Brodkorb, A. Formation of Non-Native β -Lactoglobulin during
 380 Heat-Induced Denaturation. *Food Biophysics*, 10.1007/s11483-011-9230-3

381 Kim, J.-S., & Lee, Y.-S. (2009). Study of Maillard reaction products derived from aqueous model
 382 systems with different peptide chain lengths. *Food Chemistry*, 116(4), 846-853.

383 Laemmli, U. K. (1970). Cleavage of structural proteins during assembly of head of bacteriophage-T4.
 384 *Nature*, 227(5259), 680-&.

385 Lan, X., Liu, P., Xia, S., Jia, C., Mukunzi, D., Zhang, X., Xia, W., Tian, H., & Xiao, Z. (2010).
 386 Temperature effect on the non-volatile compounds of Maillard reaction products derived from
 387 xylose-soybean peptide system: Further insights into thermal degradation and cross-linking.
 388 *Food Chemistry*, 120(4), 967-972.

389 Laroque, D., Inisan, C., Berger, C., Voulard, É., Dufossé, L., & Guérard, F. (2008). Kinetic study on
 390 the Maillard reaction. Consideration of sugar reactivity. *Food Chemistry*, 111(4), 1032-1042.

391 Ledl, F., Beck, J., Sengl, M., Osiander, H., Estendorfer, S., Severin, T., & Huber, B. (1989). Chemical
 392 pathways of the Maillard reaction. *Progress in Clinical and Biological Research*, 304, 23-42.

393 Li, Y., Lu, F., Luo, C., Chen, Z., Mao, J., Shoemaker, C., & Zhong, F. (2009). Functional properties of
 394 the Maillard reaction products of rice protein with sugar. *Food Chemistry*, 117(1), 69-74.

395 Liskova, K., Kelly, A. L., O'Brien, N., & Brodkorb, A. (2010). Effect of denaturation of alpha-
 396 lactalbumin on the formation of BAMLET (bovine alpha-lactalbumin made lethal to tumor
 397 cells). *Journal of Agricultural and Food Chemistry*, 58(7), 4421-4427.

398 Van Boekel, M. A. J. S. (1998). Effect of heating on Maillard reactions in milk. *Food Chemistry*, 62(4),
 399 403-414.

400 Medrano, A., Abirached, C., Panizzolo, L., Moyna, P., & Añón, M. C. (2009). The effect of glycation
 401 on foam and structural properties of β -lactoglobulin. *Food Chemistry*, 113(1), 127-133.

402 Miralles, B., Martínez-Rodríguez, A., Santiago, A., van de Lagemaat, J., & Heras, A. (2007). The
 403 occurrence of a Maillard-type protein-polysaccharide reaction between β -lactoglobulin and
 404 chitosan. *Food Chemistry*, 100(3), 1071-1075.

405 Morales, F. J., Romero, C., & JimenezPerez, S. (1996). Fluorescence associated with Maillard reaction
 406 in milk and milk-resembling systems. *Food Chemistry*, 57(3), 423-428.

407 Oliver, C. M., Melton, L. D., & Stanley, R. A. (2006). Creating proteins with novel functionality via
 408 the Maillard reaction: A review. *Critical Reviews in Food Science and Nutrition*, 46(4), 337-
 409 350.

410 Qi, J. R., Yang, X.-Q., & Liao, J.-S. (2009). Improvement of functional properties of acid-precipitated
 411 soy protein by the attachment of dextran through Maillard reaction. *International Journal of*
 412 *Food Science and Technology*, 44(11), 2296-2302.

413 Rojas, S. A., Goff, H. D., Senaratne, V., Dalgleish, D. G., & Flores, A. (1997). Gelation of commercial
 414 fractions of β -lactoglobulin and α -lactalbumin. *International Dairy Journal*, 7(1), 79-85.

415 Rufian-Henares, J. A., Delgado-Andrade, C., & Morales, F. J. (2006). Occurrence of acetic acid and
 416 formic acid in breakfast cereals. *Journal of the Science of Food and Agriculture*, 86(9), 1321-
 417 1327.

418 Schmitt, C., Bovay, C., & Frossard, P. (2005). Kinetics of formation and functional properties of
 419 conjugates prepared by dry-state incubation of beta-lactoglobulin/acacia gum electrostatic
 420 complexes. *Journal of Agricultural and Food Chemistry*, 53(23), 9089-9099.

421 Sun, Y., Hayakawa, S., Puangmanee, S., & Izumori, K. (2006). Chemical properties and antioxidative
 422 activity of glycated α -lactalbumin with a rare sugar, d-allose, by Maillard reaction. *Food*
 423 *Chemistry*, 95(3), 509-517.

424 Wahyuni, M., Ishizaki, S., & Tanaka, M. (1998). Improvement of thermal stability of fish water soluble
 425 proteins with glucose-6-phosphate through the Maillard reaction. *Fisheries Science*, 64(6),
 426 973-978.

427 Werner, B. (1982). Chemical changes in food by the maillard reaction. *Food Chemistry*, 9(1-2), 59-73.

428 Wooster, T. J., & Augustin, M. A. (2006). β -Lactoglobulin–dextran Maillard conjugates: Their effect
 429 on interfacial thickness and emulsion stability. *Journal of Colloid and Interface Science*,
 430 303(2), 564-572.

431 Yanagimoto, K., Lee, K. G., Ochi, H., & Shibamoto, T. (2002). Antioxidative activity of heterocyclic
432 compounds found in coffee volatiles produced by Maillard reaction. *Journal of Agricultural*
433 *and Food Chemistry*, 50(19), 5480-5484.

434 Yen, G. C., & Hsieh, P. P. (1995). Antioxidative activity and scavenging effects on active oxygen of
435 xylose-lysine Maillard reaction products. *Journal of the Science of Food and Agriculture*,
436 67(3), 415-420.

437 Yoshimura, Y., Iijima, T., Watanabe, T., & Nakazawa, H. (1997). Antioxidative effect of Maillard
438 reaction products using glucose-glycine model system. *Journal of Agricultural and Food*
439 *Chemistry*, 45(10), 4106-4109.

440

Fig. 1. Changes in pH (A) and free amino groups (B) of α -LA-Ribose and β -LG-Ribose model system after heat treatment at 95°C for up to 5 hours; distribution of native (black), non-native (white) and aggregate (grey) α -LA (C) and β -LG (D) in protein-ribose model systems during heat treatment at 95°C for up to 5 hours. Triangles represent ribose; bars indicate the standard deviation from independent triplicate measurements.

Fig. 2. (A) Fluorescence excitation (left) and fluorescence (right) spectra of α -LA-Ribose model system heated at 95°C for 0, 0.5, 1, 2, 3, 4 and 5h; Excitation wavelength for emission spectra was set to 337nm and scanning wavelength range from 370nm to 580nm. Emission wavelength for excitation spectra was 420nm and scanning wavelength range from 300nm to 400nm. Inserts (A') UV/vis absorption spectra of α -LA-Ribose (arrow corresponds to increasing heating times). (B) and Inserts (B') are the corresponding spectra for β -LG-Ribose model system. (C) shows the fluorescence intensity at 420 nm of α -LA-Ribose and β -LG-Ribose model system (excitation at 337nm).

Fig. 3. Size Distribution (by Volume) measured by dynamic light scattering of α -LA (A), α -LA-Ribose (B), β -LG (C), and β -LG-Ribose (D) during heat treatment of 95°C for up to 5 hours with an initial pH 8.4.

Fig. 4. SDS-PAGE of 3% (w/w) α -LA (A and B) and β -LG (C and D) in the presence of 3% (w/w) ribose treated at 95°C for various times. (A) and (C) represent SDS-PAGE under non-reducing and (B) and (D) under reducing (presence of β -mercaptoethanol) conditions. Lanes 1-7 correspond to heating times of 0, 0.5, 1, 2, 3, 4 and 5 hours; the far right lane corresponds to molecular weight markers.

Fig. 5. Size exclusion chromatograms of (A) α -LA-Ribose and (B) β -LG-Ribose system during heating (95°C) for different times (from 0 to 5h) at pH 8.4.

Fig. 6. Antioxidant activity of the α -LA-Ribose and β -LG-Ribose during heating (95°C) for different times (from 0 to 5h) at pH 8.4. (A) DPPH radical scavenging activity and (B) relative reducing power.

Fig. 1. Changes in pH (A) and free amino groups (B) of α -LA-Ribose and β -LG-Ribose model system after heat treatment at 95°C for up to 5 hours; distribution of native (black), non-native (white) and aggregate (grey) α -LA (C) and β -LG (D) in protein-ribose model system during heat treatment at 95°C for up to 5 hours. Triangles represent ribose; bars indicate the standard deviation from independent triplicate measurements.

475

476

477

478

480

481 Fig. 2. (A) Fluorescence excitation (left) and fluorescence (right) spectra of α -LA-Ribose model
 482 system heated at 95°C for of 0, 0.5, 1, 2, 3, 4 and 5h; Excitation wavelength for emission spectra was
 483 set to 337nm and scanning wavelength range from 370nm to 580nm. Emission wavelength for
 484 excitation spectra was 420nm and scanning wavelength range from 300nm to 400nm. Inserts (A')
 485 UV/vis absorption spectra of α -LA-Ribose (arrow corresponds to increasing heating times). (B) and
 486 Inserts (B') are the corresponding spectra for β -LG-Ribose model system. (C) shows the fluorescence
 487 intensity at 420 nm of α -LA-Ribose and β -LG -Ribose model system (excitation at 337nm).

488

489

490

491

492

493 Fig. 3. Size Distribution (by Volume) measured by dynamic light scattering of α -LA (A), α -LA-Ribose
494 (B), β -LG (C), and β -LG-Ribose (D) during heat treatment of 95°C for up to 5 hours with an initial pH
495 8.4.
496

498

499 Fig. 4. SDS-PAGE of 3% (w/w) α -LA (A and B) and β -LG (C and D) in the presence of 3% (w/w)
 500 ribose treated at 95°C for various times. (A) and (C) represent SDS-PAGE under non-reducing and (B)
 501 and (D) under reducing (presence of β -mercaptoethanol) conditions. Lanes 1-7 correspond to heating
 502 times of 0, 0.5, 1, 2, 3, 4 and 5 hours; the far right lane corresponds to molecular weight markers.

503

504

505

506

507 Fig. 5. Size exclusion chromatograms of (A) α -LA-Ribose and (B) β -LG-Ribose system during heating
 508 (95°C) for different times (from 0 to 5h) at pH 8.4.

509

Fig. 6. Antioxidant activity of the α -LA-Ribose and β -LG-Ribose during heating (95°C) for different times (from 0 to 5h) at pH 8.4. (A) DPPH radical scavenging activity and (B) relative reducing power.