**Variation in bioactive content in broccoli (*Brassica oleracea*
var. *italica*) grown under conventional and organic
production systems**

Journal:	<i>Journal of the Science of Food and Agriculture</i>
Manuscript ID:	JSFA-14-0882.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Valverde, Juan; Monaghan Mushrooms, Research Reilly, Kim; Teagasc, Food Research Centre Villacreces, Salvador; Teagasc, Food Research Centre Gaffney, Michael; Teagasc, Food Research Centre Grant, Jim; Teagasc, Food Research Centre Brunton, Nigel; University College Dublin, School of Agriculture and Food Science
Key Words:	<i>Brassica oleracea</i> , organic agriculture, glucosinolates, neoglucobrassicin, glucobrassicin, phenolic compounds

SCHOLARONE™
Manuscripts

Variation in bioactive content in broccoli (*Brassica oleracea* var. *italica*) grown under conventional and organic production systems

Running title: organic_broccoli_JSFA

Juan Valverde,^{*1,2} Kim Reilly,^{*1} Salvador Villacreces,¹ Michael Gaffney,¹ James Grant,¹ and Nigel Brunton^{1,3}

*** Both authors contributed equally**

- ¹ Teagasc, Ashtown Food Research Centre, Ashtown, Dublin 15, Ireland.
- ² Monaghan Mushrooms, Tyholland, Co. Monaghan, Ireland.
- ³ School of Agriculture and Food Science, University College Dublin, Ireland.

Correspondence: Dr Kim Reilly, Teagasc Food Research Centre, Ashtown, Dublin 15, Ireland. Tel: +353 (0) 18 05 99 50, e-mail: kim.reilly@teagasc.ie

Abstract

BACKGROUND: Broccoli and other cruciferous vegetables contain a number of bioactive compounds - in particular glucosinolates and polyphenols, which are proposed to confer health benefits to the consumer. Demand for organic crops is at least partly based on a perception that organic crops may contain higher levels of bioactive compounds; however insufficient research has been carried out to either support or refute such claims.

RESULTS: In this study we examined the effect of conventional, organic, and mixed cultivation practices on the content of total phenolics, total flavonoids, and total and individual glucosinolates in two varieties of broccoli grown over two years in a split-plot factorial systems comparison trial. Levels of total phenolics and total flavonoids showed a significant year on year variation but were not significantly different between organic and conventional production systems. In contrast, levels of the indolyl glucosinolates glucobrassicin and neoglucobrassicin were significantly higher ($p < 0.05$) under fully organic compared to fully conventional management.

CONCLUSION: Organic cultivation practices resulted in significantly higher levels of glucobrassicin and neoglucobrassicin in broccoli florets, however other investigated compounds were unaffected by production practices.

Keywords: *Brassica oleracea*; organic agriculture; glucosinolates; neoglucobrassicin; glucobrassicin; phenolic compounds.

51 **INTRODUCTION**

52 Over the last 10 years demand for organically produced food products has increased and this
53 has been recognized as an important consumer trend in the USA and Europe.¹ Although the
54 reasons behind this increasing demand differ among countries and consumer groups, the
55 perception of improved animal welfare, environmental protection, and food quality
56 characteristics (including health, nutritional and/or sensory attributes) are among some of the
57 reasons cited for this trend.^{1,2} Organic crop production in the European Union is carried out
58 under strictly defined production practices.^{3,4} Despite consumer presumption that organic
59 fruits and vegetables are healthier, the reality is that previous studies on nutrient and
60 phytochemical composition have shown contradictory results.⁵⁻¹² One of the main challenges
61 for these types of studies is the design of a robust experiment that takes account of important
62 factors which can contribute to variability between studies such as crop variety, geographical
63 location, growth season, management practices used, experimental design and statistical
64 power.

65 Levels of glucosinolates and phenolic compounds in broccoli appear to be affected
66 primarily by variety although only a few studies have been carried out with a number of
67 cultivars examined under uniform cultivation conditions.¹³⁻¹⁷ Levels are also affected by
68 factors such as nitrogen fertilization, environmental factors and season,¹⁸⁻² although levels of
69 both phenolic compounds and glucosinolates appear to remain stable under post-harvest
70 storage treatments designed to simulate commercial storage and marketing.²²

71 Broadly three types of study have been used to examine differences in nutritional or
72 phytochemical content between organic and conventional foods. The first type comprise
73 basket surveys, in which sampling is made at retail points, by grouping samples according to
74 production system e.g.⁹ The second type of study uses matched paired farms where existing
75 farms using either conventional or organic production systems are matched as far as possible

1
2
3 76 in terms of location and crop produced.²³ The last type of study is by performing replicated
4
5 77 field trials.²⁴ In any of these types of study a number of uncontrolled factors can influence
6
7 78 the results obtained. Consequently, contradictory results have been obtained between different
8
9 79 studies. In an extensive and widely reported meta-analysis Dangour *et al.*⁵ note the
10
11 80 difficulties and variable quality of research in the area and suggest five criteria for a study of
12
13 81 acceptable quality: firstly to clearly define the organic production methods including the name
14
15 82 of the certification body; secondly to specify crop cultivar or animal breed; thirdly to state
16
17 83 which nutritionally relevant substance was analysed; fourthly to state methods of analysis and
18
19 84 lastly to state methods used for statistical analyses.
20
21
22

23 85 The aim of this study was to compare the content of glucosinolates and phenolic
24
25 86 compounds in organically and conventionally grown broccoli (*Brassica oleracea* var. *italica*).
26
27
28

29 88 **MATERIALS AND METHODS**

30 89 **Experimental design**

31
32
33 90 Two varieties each of broccoli, carrot and onion were grown in a replicated split-plot systems
34
35 91 comparison trial. The field trial has a factorial design which divides both organic and
36
37 92 conventional production systems into component soil management and pest-control practices.
38
39 93 It was designed to investigate the effect of, and any interaction between, production system
40
41 94 components - a) soil management and b) pest-control measures, that differ between organic
42
43 95 and conventional systems. In order to encompass seasonal variation, sampling and data
44
45 96 analysis presented is on two years of crop production. The trial design and statistical analyses
46
47 97 comply with the suggested quality criteria of Dangour *et al.*⁵ The trial design includes fully
48
49 98 organic, fully conventional and mixed treatments, which allows comparison of fully organic
50
51 99 and fully conventional production, whilst allowing investigation of soil and pest-control
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

components which make up organic or conventional agricultural practices. Management practices used are summarised in Table 1.

Plant material

Broccoli samples for this study were grown in 2009 (year1) and 2010 (year 2). Varieties selected were cv. ‘Belstar’ and cv. ‘Fiesta’ both of which are commercial varieties commonly grown by conventional and organic growers in Ireland. Plants were grown according to cultivation practices in compliance with European and Irish standards for organic certification (Irish Organic Farmers and Growers Association (IOFGA) and Organic Trust standards, see details below) and/or following the Irish Agriculture and Food Development Authority (Teagasc) recommendations for conventional practices.^{25, 26} The year 1 broccoli crop was sown on 24th April, transplanted on 25th June and harvested between 4th and 25th September 2009. The year 2 crop was sown on 23rd March, transplanted on 18th May and harvested between 27th July and 5th August 2010. Plants were produced as modular transplants in 216 trays and transplanted at 40 cm in-row spacing with 2 rows per 1.52 m bed. External rows were treated as guard rows with samples harvested from internal rows only. Specific applied inputs for broccoli propagation and cultivation are shown in Table 2. Climatic conditions during the growing season in both years are shown in supplementary material (Table S1). Additional information on the field trial is available at <http://www.ipfn.ie/publications/agronomic/>.

For each experimental plot broccoli was harvested at commercial maturity from the internal rows with guard rows excluded. The mean floret weight was calculated as the total weight of harvested broccoli florets divided by the number of florets. Samples for analysis were primary florets of similar size and were immediately refrigerated and then frozen at -20°C within 24 hours of harvest. Samples from each experimental plot were composite samples comprising three healthy, disease free florets of marketable quality.

125

Field trial

The field trial is located at Teagasc Kinsealy Research Centre, Kinsealy (53° 25' N Lat 6° 10' W), in north county Dublin, Ireland. Soil type was characterised as loam to clay loam belonging to the grey brown podzolic soil group and with a high base status (Altitude: 28 metres O.D. (ordnance datum), Slope: 1° Drainage: Moderately well drained). Soil type was consistent across the experimental trial site as indicated by a previous detailed soil map of the area.²⁷ Equivalent rates of nitrogen (N), phosphorus (P) and potassium (K) were applied to both conventional and organic soil treatments for each crop and the rates applied were according to Teagasc published recommendations for the crop.²⁵ Although the amount of N, P and K supplied was identical between systems, plant availability and uptake is affected by fertilizer form (chemical vs. organic) due to differences in water solubility and the need for organic fertilizer to be broken down by soil microbes. The trial was established in spring 2009 on land which had previously been under grass set-aside for over 10 years. The same irrigation source was used for the whole trial.

The trial was a 2 x 2 x 2 factorial split-plot design, with 4 replicates (blocks). Variety was assigned as the main plot, with pest-control and soil treatment assigned as sub plots. Two varieties (V1, V2) of 3 crops (carrots, broccoli and onion) were grown in each year. There were two levels of soil treatment – an organic soil treatment (OS) and a conventional soil treatment (CS); and 2 levels of pest control – an organic pest-control treatment (OP) and a conventional pest-control treatment (CP).

The OS treatment comprised certified organic fertilizer inputs, a 4 year horticultural crop rotation including a red clover ley (*Trifolium repens*) and use of winter cover crops (Table 1). The CS treatment comprised use of mineral fertilizers, with no set crop rotation. Equivalent rates of nitrogen (N), phosphorus (P) and potassium (K) were applied to both CS

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

and OS treatments for each crop following a spring soil test and rates were according to Teagasc recommendations for the crop.²⁵ Fertilizer was applied as calcium ammonium nitrate (CAN), single super-phosphate and sulphate of potash for the CS treatment, or Greenvale (3:3:1) and ProKali (3:0:14) for the OS treatment. Specific inputs are shown in Table 2.

Pest-control measures were required for control of weeds, cabbage root-fly (*Delia radicum*), and caterpillars of Diamond-back moth (*Plutella xylostella*) and Large White butterfly (*Pieris brassicae*). Organic pest-control (OP) measures comprised mechanical (e.g. hand hoeing, brassica collars) and certified organic treatments as shown in Table 2. Conventional pest-control (CP) treatments involved a chemical spay programme and were in accordance with an Integrated Pest Management plan, in keeping with commercial growing practises in North Dublin.

Within each replicate (n=4) each crop is grown under 8 possible treatment combinations (V1+OS+OP, V1+OS+CP, V1+CS+OP, V1+CS+CP, V2+OS+OP, V2+OS+CP, V2+CS+OP, V2+CS+CP) giving a total of 32 plots per crop per year (Supplementary material Figures S1, S2). The organic cultivation practices used were in compliance with EC 834/2007³ and with national standards for organic certification set out by the Irish organic certification bodies (IOFGA and the Irish Organic Trust) with the exception that for experimental purposes the separation distance (generally 50 m) required between adjacent organic and conventional enterprises was not practised between organic and conventional treatment plots.

Each experimental plot comprised two 1.52 m beds of 5.5 m length (area 16.7 m²). In order to prevent or reduce as far as possible any cross contamination between treatment plots, within each replicate each plot was separated from neighbouring treatment plots by a 1 m wide untreated grass inter-plot area, with 3 m grass areas between replicate blocks. Fertilizers were applied by hand and tractor tillage operations carried out such as to minimise movement

1
2
3 175 of soil between treatments. Pesticides were applied using a knapsack sprayer with hood to
4
5 176 prevent spray drift between treatment plots. Although it was not practically feasible to
6
7 177 completely prevent all movement between plots (e.g. arthropods, earthworms, microbes could
8
9 178 be expected to move through the soil to some extent) statistically significant differences in
10
11 179 soil microbial activity and functional diversity between plots under different management
12
13 180 practices has been demonstrated in this trial ²⁸ indicating that measures used were effective in
14
15 181 allowing different soil biology in different plots.
16
17

18
19 182 Diagrams showing the plot allocation and crop rotation in years 1 and 2 are shown in
20
21 183 supplementary material Figures S1 and S2.
22
23 184

24 25 185 **Total phenolics and total flavonoids**

26
27 186 For the determination of total phenolics a modification of the Folin-Ciocalteu method was
28
29 187 used. Briefly, broccoli samples were ground to a fine powder under liquid nitrogen. Frozen
30
31 188 tissue (0.50 g) was transferred to a falcon tube and 5 mL of 80 % methanol (v/v) was added.
32
33 189 Tubes were vortexed thoroughly and allowed stand at room temperature for 20 minutes.
34
35 190 Tubes were mixed by inversion and 1.5 mL aliquots of extract were transferred to a microfuge
36
37 191 tube. Microfuge tubes were centrifuged at 12,000g for 5 minutes at 4 °C and the supernatant
38
39 192 was transferred to a fresh tube. For each sample 150 µL of the methanolic extract, 150 µL 80
40
41 193 % (v/v) methanol, 150 µL Folin-Ciocalteu reagent and 1050 µL sodium carbonate solution
42
43 194 (20 % w/v) were pipetted into a microfuge tube, vortexed and placed in the dark at room
44
45 195 temperature for 20 minutes. Tubes were centrifuged at 12,000g for 3 minutes and the
46
47 196 supernatant was transferred to a fresh tube. The absorbance at 725 nm (A_{725}) was determined
48
49 197 relative to a blank containing 80% (v/v) methanol instead of extract, and the concentration
50
51 198 was determined from a calibration curve using gallic acid. Results are expressed as gallic acid
52
53 199 equivalents on a fresh weight basis (GAE mg/100g FW).
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

200 Determination of flavonoids was according to Marinova *et al.*²⁹ For each sample 150
201 μL of the methanolic extract and 600 μL MilliQ water were added to a microfuge tube and
202 mixed by inversion. To each tube 45 μL of 5% sodium nitrite (NaNO_2) was added and tubes
203 were incubated at room temperature for 5 minutes. To each tube 45 μL of 10% aluminium
204 chloride was added and tubes incubated for a further 1 minute. Subsequently 300 μL of 1M
205 sodium hydroxide (NaOH) and 360 μL MilliQ water was added and tubes mixed vigorously.
206 The absorbance at 510 nm (A_{510}) was measured relative to a blank containing 80% (v/v)
207 methanol instead of extract, and flavonoid concentration was determined from a standard
208 curve using catechin as a standard. Results are expressed as catechin equivalents (CE
209 $\text{mg}/100\text{g}$ FW).

210
211 **Glucosinolate extraction**

212 Frozen broccoli samples were freeze dried in a large scale freeze drier (*Frozen in Time Ltd.*
213 United Kingdom). Once freeze dried, samples were milled, vacuum packed in polypropylene
214 bags and kept at -80°C until analysis. Glucosinolates were extracted from freeze-dried
215 broccoli powder using pressurised liquid extraction with an ASE 200 instrument (Dionex;
216 Sunnyvale, CA, USA) with attached solvent controller. Glucosinolate extraction was carried
217 out in 22 mL steel cartridges packed with a mixture of freeze-dried broccoli (1.00 g) and
218 technical grade silica to disperse sample. Extraction conditions were as described in
219 Hernandez-Hierro *et al.*³⁰

220 **Sulphatase extraction procedure**

221 Sulphatase (Type H-1 from *Helix pomatia*, Sigma, MO, USA) was purified by dissolving the
222 sulphatase powder (70 mg) in deionised water (3 mL) and adding ethanol (3 mL). This
223 solution was centrifuged (12000 rpm, 10 min, room temperature) and to the supernatant
224 ethanol (9 mL) was added after which the solution was centrifuged (12000 rpm, 10 min, room

temperature) again. The pellet was dissolved in deionised water (2 mL) and this sulphatase solution was subsequently passed through a 0.5 mL DEAE Sephadex A-25 and a 0.5 mL SP Sephadex C-25 column. This solution was collected in a vial and kept at -80 °C until use.

An aliquot (1mL) of glucosinolate extract was applied to a DEAE Sephadex A-25 column (0.5mL) and the unbound material was removed by washing with deionised water (2 x 1mL) and sodium acetate buffer (2 x 0.5mL, 20 mM, pH 5.0). After washing, purified sulfatase solution prepared as above was added and the columns were incubated overnight at room temperature. After overnight incubation, the desulphoglucosinolates (dGLS) were eluted from the columns with deionised water (3 x 1ml). The collected eluate was dried under constant N₂ flow and re-dissolved in deionised water (200µL)

Micellar Electrokinetic Capillary Chromatography

Analyses were performed using a CE capillary electrophoresis system (Agilent, Waldbronn, Germany) equipped with diode array detector. All separations were performed on a fused silica capillary (Agilent, Stevens Creek, CA; 75 µm ID, 64.5 cm total length, 56 cm effective length). Samples were injected from the anodic end of the capillary (vacuum injection, 50 mbar, 1 s). The separation buffer consisted of sodium chlorate (250 mM) and boric acid (200 mM) at pH 8.5; the separation was carried out at 12 kV and 60 °C. The capillary was conditioned between each run sequentially with 1.0 M NaOH (3 min), 0.1 M NaOH (1 min), water (1 min) and separation buffer (5 min). Detection was performed on column at 230 and 280 nm. Data processing was carried out with Chemstation software (Agilent Waldbronn, Germany). The quantity of desulfo-glucosinolates was estimated as the average of quantities calculated by comparison of normalized area under the curve (area under the curve divided by migration time) of each identified desulfo-glucosinolate peak with the normalized area under the curve of glucotropaeolin (internal standard, from the laboratory collection). Identification

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

of individual desulfo-glucosinolates was performed by calculating their migration time relative to the migration time of glucotropaeolin and their PDA profile

Statistical analysis

Statistical analysis was carried out using SAS 9.1 (Cary, NC). Floret weight, total phenolic, total flavonoid and glucosinolate data were analysed using an ANOVA mixed model containing a contrast code to compare the fully organic (OS+OP) and fully conventional (CS+CP) treatments as well the individual treatments and interactions (SAS 9.1). Pearson correlation coefficients were calculated between total phenolics, flavonoids and mean floret weights using SAS 9.1.

RESULTS AND DISCUSSION

Yield and quality

In crop production the main differences between organic and conventional growing systems involve the use of organic manures and crop rotations instead of inorganic fertilizers; and mechanical or biological methods (including naturally derived compounds) for pest-control instead of synthetic pesticides. Thus it can be considered that organic and conventional agriculture differ in two major respects – how the soil fertility is managed, and how pests are managed. Both factors may impact on crop yield and quality.

Analysis of broccoli yield, total phenolic content and total flavonoid content is shown in Table 3. Year showed a significant ($p<0.01$) effect with some measures showing a year x treatment interaction therefore data for each year was analysed separately (Table 3). Floret weight, quality and yield was higher in year 2 than in year 1 for both varieties. Mean floret weight ranged from 230.6 ± 20.3 g to 307.0 ± 17.0 g in year 1 and was considerably higher at 247.1 ± 78.5 g to 449.2 ± 19.9 g in year 2. Climate data for the trial site (Supplementary

material, Table S1) shows the growing season in year 1 was overall slightly warmer but much wetter than year 2 with excessive rainfall during the summer especially in July. In year 1 a significant soil treatment effect on floret weight ($p<0.01$) was found, with crops grown with the conventional soil (CS) treatment showing higher floret weights, however this effect was not significant in year 2. This may be due to soil nutrient leaching and/or poor root establishment in year 1 as a result of heavy rainfall. Since nutrients in conventionally fertilized soil are more readily crop available, it is possible that rainfall effects were exacerbated in the organically fertilized soil. Year 1 was a poor year for crop growth with frequent heavy rain and this was reflected in lower mean floret weight across all treatments. Year 2 was an improved year for crop production with a better yield seen for both varieties. Variety 'Belstar' performed particularly well under improved climatic conditions with a more typical level of rainfall in year 2 as indicated by the significant main effect of variety ($p=0.0242$) in this year.

In both years infestations with caterpillars of Large White butterfly and Diamond-back moth occurred throughout the crop and were effectively treated with Pyrethrum in the organic pest-control (OP) plots and with Decis in the conventional pest-control (CP) plots. At harvest broccoli florets were scored for incidence of insect or other damage, and fungal and bacterial diseases prevalent in North Country Dublin including White Blister (*Albugo candida*) and Wet Rot (*Erwinia carotovora* and *Pseudomonas* spp.). Levels of insect damage and disease were generally low and were not different between treatments in either year.

Overall no significant differences in yield were found between fully organic (OS+OP) and fully conventional (CS+CP) management in any year (Table 3). However a significant soil treatment main effect was found for floret weight in year 1 only with lower weights recorded for broccoli grown under organic soil management (OS). These data indicate that soil treatment rather than pest-control was the primary driver of broccoli yield and quality. We

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

suggest that under favourable conditions such as year 2, both organic and conventional fertilizer performed well. But under stressful conditions (such as year 1) the conventional soil (CS) treatment provided readily plant available mineral fertilizers which enabled these plants to outperform those under organic soil (OS) treatment.

Total phenolic and total flavonoid content

Levels of total phenolics (Figure 1 and Table 3) in year 1 were in the range 267.0 ±18.4 GAE mg/100g FW to 376.5±19.0 GAE mg/100g FW, and were considerably lower in year 2, ranging from 69.3±11.5 to 106.8±7.5 GAE mg/100g FW. Thus levels in year 2 equate to around one third to one quarter of the levels found in year 1. Total flavonoid content showed the reverse trend (Figure 2 and Table 3) and ranged from 5.6±1.1 to 25.9±3.4 CE mg/100 FW in year 1, with higher levels in year 2 ranging from 23.8±3.5 to 35.4±5.2 CE mg/100 FW across treatments.

The major phenolic compounds found in broccoli include flavonols such as quercetin and kaempferol glycosides, hydroxycinnamoyl derivatives and chlorogenic acids.²² Phenylalanine ammonia lyase (PAL) the key entry point enzyme for synthesis of phenolic compounds is well known to be up-regulated by stresses including UV light, low temperature, nutrient deficiency, wounding and pest or pathogen attack.³¹ The branch pathway to flavonoid synthesis is controlled by chalcone synthase (CHS). Flavonoid synthesis is commonly reported to be up-regulated by light in several crops including broccoli. In a three year field study which examined levels of the flavonols kaempferol and quercetin in three broccoli varieties (‘Marathon’, ‘Lord’ and ‘Fiesta’) the level of total solar radiation over the growing period had a significant effect on both flavonols with higher levels under increased radiation.²⁰ We propose that the higher levels of total flavonoids in year 2 in this study are due to higher light levels; whilst the high overall content of total phenolics in year 1 reflect increased

323 production of phenolic acids in response to stress caused by heavy rainfall and associated
324 waterlogging of soils.

325 As with yield no significant differences in total phenolic or flavonoid content were
326 found between fully organic (OS+OP) and fully conventional (CS+CP) management in any
327 year (Table 3). However a significant soil treatment main effect was found for total phenolics
328 and total flavonoids in year 1 only. Data indicated that in a poor year for crop growth (year 1)
329 broccoli floret total phenolic and flavonoid content was increased when crops were grown in
330 organic soil (OS). We ascribe this result to nutrient stress of the plants grown in the OS
331 treatment. There was a significant strong positive correlation ($p < 0.001$) in both years between
332 total phenolic content and flavonoid content with $r = 0.74$ in year 1 and $r = 0.69$ in year 2.

333 The levels of total phenolics and flavonoids reported here are in agreement with levels
334 found in green broccoli varieties in other studies.¹³⁻¹⁶ Relatively few studies have compared
335 phenolic content in vegetable crops grown under conventional and organic systems and to our
336 knowledge no previous field trial studies have examined phenolic content in organic and
337 conventionally grown broccoli. A recent well controlled Danish study⁷ measured levels of
338 flavonoids in onions and phenolic acids in carrots and potatoes grown over two years using
339 either conventional or two types of organic system (cover crop fertility building or animal
340 manure based fertilizer regime). The predominant phenolic acid in potatoes (5-caffeoylquinic
341 acid) was significantly higher in the cover crop based organic system than in the conventional
342 system.⁷ In contrast phenolic acids in carrot and flavonoids in onion showed a large year to
343 year variation but were not affected by production system. Our data indicate that total
344 phenolic and flavonoid content in broccoli shows significant year to year variation but is not
345 significantly different in fully organic (OS+OP) compared to fully conventional (CS+CP)
346 production systems.

347

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Total and individual glucosinolate content

Glucosinolate contents determined in broccoli samples are shown in Table 4. Peaks from all samples were resolved and easy to identify (Figure 3). The desulfoglucosinolates from broccoli were characterised by the presence of aliphatic glucosinolates and indol-3-yl glucosinolates. Mainly aliphatic glucosinolates - glucoraphanin, glucobrassicin, and neoglucobrassicin were the major components in most samples. 4-MeO-glucobrassicin was found at lower levels and progoitrin and sinigrin were found inconsistently in trace amounts in some samples. No evidence of glucoiberin nor 4-OH-glucobrassicin was found. Total glucosinolate content ranged from 3.0 to 20.9 $\mu\text{mol/g DW}$ and averages were in the range 8.1 $\pm$ 0.7 to 10.1 $\pm$ 0.4 $\mu\text{mol/g DW}$ ($\pm$ standard error, $n = 4$) depending on treatment. Previous studies have shown that either glucoraphanin or glucobrassicin is the predominant glucosinolate in broccoli.^{13-20,32-35} It is not yet well understood why in some studies one is in higher proportion than the other and this may be due to genetic (i.e. differences between cultivars) and/or environmental conditions (for review see ³⁶). In this study, the average concentration ($\pm$ standard error, $n = 4$) of major glucosinolates across production systems and over the two year period ranged from 1.8 $\pm$ 0.2 to 3.3 $\pm$ 0.3 $\mu\text{mol/g DW}$ for glucoraphanin, 2.9 $\pm$ 0.2 to 4.7 $\pm$ 0.3 $\mu\text{mol/g DW}$ for glucobrassicin, 0.2 $\pm$ 0.02 to 0.4 $\pm$ 0.03 $\mu\text{mol/g DM}$ for 4-MeO-glucobrassicin and 1.4 $\pm$ 0.2 to 3.1 $\pm$ 0.1 $\mu\text{mol/g DW}$ for neoglucobrassicin respectively. This is in agreement with ranges of glucosinolates in broccoli previously reported.^{13, 17} No significant interaction or main effect of year on total or individual glucosinolate content was observed across the two year period. This is important because a significant difference in the content of secondary metabolites between years would indicate significant seasonal environmental interactions in the glucosinolate profile. Variability in biochemical data between years is often observed and is normally considered to be due to the crops response to different climatic conditions.³²⁻³⁴ Differences in broccoli glucosinolate content due to

environmental conditions, in particular temperature and irradiation levels, have been reported in other studies.^{20, 33} In the two seasons reported here, mean temperatures, humidity and wind speed were similar in both years, but rainfall levels were almost double in year 1 relative to year 2. Therefore rainfall levels do not appear to impact glucosinolate content. A number of previous studies have indicated a significant genotype effect on glucosinolate profile in broccoli.^{13,17,34,35} Our data indicated levels of sinigrin were significantly higher ($p<0.05$) in cv. 'Fiesta' than in cv. 'Belstar' across treatments and years. Conversely glucoraphanin and 4-MeO-glucobrassicin were significantly higher ($p<0.05$) in cv. 'Belstar' than in cv. 'Fiesta' across treatments and years (Figure 4 and Table 4). Glucoraphanin has been extensively studied due to the potential bioactivity of its isothiocyanate breakdown product sulforaphane.³⁷ The finding of consistently higher levels in 'Belstar' is therefore of relevance from a health perspective. Levels of 4-MeO-glucobrassicin were lower in both varieties and its potential as bioactive compound has been less explored.

Glucobrassicin and neoglucobrassicin content was significantly ($p<0.05$) higher in samples grown under fully organic treatment (organic soil and organic pest-control; OS+OP) compared to samples grown under completely conventional treatment (CS+CP). Mixed model ANOVA showed that no significant main or interaction effects were observed for different soil treatments (Table 4). For neoglucobrassicin a significant variety by pest-control interaction was seen. It is important to highlight that although glucobrassicin contents were not significantly different for pest-control, values were near the 95% confidence threshold ($p=0.067$).

In terms of potential bioactivity the breakdown product of indolyl glucosinolates including glucobrassicin and neoglucobrassicin is indole-3-carbinol (I3C). I3C and its condensation product 3,3' diindolylmethane (DIM) have recently received considerable attention as anti-carcinogenic compounds. They exhibit potent anti-tumor activity with low levels of toxicity

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

in a wide range of human cancer cell lines.^{38,39} The finding of statistically significant higher levels of glucobrassicin and neoglucobrassicin in broccoli grown under organic management practices would appear to be especially robust since similar results were obtained in an earlier basket study.⁹ In the study of Meyer *et al.*⁹ levels of individual glucosinolates were profiled in market purchased broccoli obtained at monthly intervals over a 1 year period in Germany. Results indicated no significant differences in glucoraphanin content, whilst neoglucobrassicin and glucobrassicin were significantly higher ($p<0.01$) in organic than in conventional broccoli. The finding of similar results in two different types of study (market study and field trial) strengthen the conclusion that the indolyl glucosinolates glucobrassicin and neoglucobrassicin consistently occur at higher levels in organically grown broccoli across different varieties and geographical locations.

CONCLUSIONS

Our data indicated that yield (mean floret weight), total phenolic and flavonoid content in broccoli shows significant year on year variation, but is not significantly different in organic (OS+OP) compared to conventional (CS+CP) production systems. We hypothesise that in year 1 increased stress caused a generalised increase in total phenolic content via up-regulation of PAL – the key enzyme controlling entry of metabolites into central phenylpropanoid metabolism. We further hypothesise that since in year 2 environmental conditions were more favourable, PAL was not up-regulated, however the chalcone synthase controlled branch pathway into flavonoid synthesis, would be up-regulated by light. Thus under these conditions a greater proportion of phenolic synthesis would be shunted towards flavonoid synthesis. Further studies to investigate this hypothesis would be of interest.

In contrast total and individual glucosinolate content was unaffected by season, but levels of two specific glucosinolates – glucobrassicin and neoglucobrassicin - were

significantly higher in the fully organic production system. Levels of glucorophanin were also consistently higher in variety 'Belstar' than in 'Fiesta'. These data underscore the nuanced regulation of levels of bioactive compounds in crop plants. It is becoming clear that plant foods contain a wide diversity of bioactive compounds which may be affected by genotype, and also respond differently to the plant's environment depending on the specific metabolite involved. We suggest that specific bioactive compounds will be responsive to production practices used in either organic or conventional agriculture whilst others will not. This complexity may account for much of the variability seen in literature reviews and meta-analyses (e.g. ^{5, 40, 41}) comparing organic and conventional food, and highlights the need for further well designed studies focussed on specific metabolites or groups of metabolites in known crop varieties.

ACKNOWLEDGEMENTS

The authors thank Prof. Carlo Leifert, University of Newcastle (UK), Dr. Charles Merfield, Christy Roberts and Leo Finn (Teagasc) for assistance in designing, implementing and maintaining the field trial, and Prof. Hilmer Sørensen and Jens C. Sørensen, University of Copenhagen (Denmark) for their help with MECC. The Irish Department of Agriculture, Fisheries and Food (FIRM 06/NITAFRC6) is gratefully acknowledged for financial support of this work.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Falguera V, Aliguer N, Falguera M, An integrated approach to current trends in food consumption: Moving toward functional and organic products? *Food Control* **26(2)**: 274-281 (2012).

2. Janssen M, Hamm U, Product labelling in the market for organic food: Consumer preferences and willingness-to-pay for different organic certification logos. *Food Quality and Preference* **25(1)**: 9-22 (2012).

3. EU, Council Regulation (EC) No 834/2007 of 28th June 2007 on organic production and labelling of organic products and repealing Regulation (EEC) No 2092/91 Official Journal of the European Union, 20.7.2007.(2007).

4. EU, Council Regulation No. 2092/91 of 24th June on organic production of agricultural products and indications referring thereto on agricultural products and foodstuffs. OJL 198 22.7.p.1 1991. (1991).

5. Dangour AD, Dodhia SK, Hayter A, Allen E, Lock K, Uauy R, Nutritional quality of organic foods: a systematic review. *American Journal of Clinical Nutrition* **90(3)**: 680-685 (2009).

6. Soltoft M, Eriksen MR, Trager AWB, Nielsen J, Laursen KH, Husted S, Halekoh U, Knuthsen P, Comparison of Polyacetylene Content in Organically and Conventionally Grown Carrots Using a Fast Ultrasonic Liquid Extraction Method. *Journal of Agricultural and Food Chemistry* **58(13)**: 7673-7679 (2010).

7. Soltoft M, Nielsen J, Lauren KH, Husted S, Halekoh U, Knuthsen P, Effects of Organic and Conventional Growth Systems on the Content of Flavonoids in Onions and Phenolic Acids in Carrots and Potatoes. *Journal of Agricultural and Food Chemistry* **58(19)**: 10323-10329 (2010).

- 468 8. Reganold JP, Andrews PK, Reeve JR, Carpenter-Boggs L, Schadt CW, Alldredge JR,
469 Ross CF, Davies NM, Zhou JZ, Fruit and Soil Quality of Organic and Conventional
470 Strawberry Agroecosystems. *Plos One* **5(9)**: e12346. doi:10.1371/journal.pone.0012346
471 (2010).
- 472 9. Meyer M, Adam ST, Comparison of glucosinolate levels in commercial broccoli and
473 red cabbage from conventional and ecological farming. *European Food Research and*
474 *Technology* **226(6)**: 1429-1437 (2008).
- 475 10. Chen P, Harnly JM, Lester GE, Flow Injection Mass Spectral Fingerprints
476 Demonstrate Chemical Differences in Rio Red Grapefruit with Respect to Year, Harvest
477 Time, and Conventional versus Organic Farming. *Journal of Agricultural and Food*
478 *Chemistry* **58(8)**: 4545-4553 (2010).
- 479 11. Magkos F, Arvaniti F, Zampelas A, Organic food: Buying more safety or just peace of
480 mind? A critical review of the literature. *Critical Reviews in Food Science and Nutrition* **46**
481 **(1)**: 23-56 (2006).
- 482 12 Rembialkowska E, Quality of plant products from organic agriculture. *Journal of the*
483 *Science of Food and Agriculture* **87**: 2757-2762 (2007).
- 484 13. Vallejo F, Tomas-Barberan FA, Garcia-Viguera C, Potential bioactive compounds in
485 health promotion from broccoli cultivars grown in Spain. *Journal of the Science of Food and*
486 *Agriculture* **82(11)**: 1293-1297 (2002).
- 487 14. Gliszczynska-Swiglo A, Kaluzewicz A, Lemanska K, Knaflewski M, Tyrakowska B,
488 The effect of solar radiation on the flavonol content in broccoli inflorescence. *Food Chemistry*
489 **100(1)**: 241-245 (2007).
- 490 15. Robbins RJ, Keck AS, Banuelos G, Finley JW, Cultivation conditions and selenium
491 fertilization alter the phenolic profile, glucosinolate, and sulforaphane content of broccoli.
492 *Journal of Medicinal Food* **8(2)**: 204-214 (2005).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

16. Vallejo F, Garcia-Viguera C, Tomas-Barberan FA, Changes in broccoli (*Brassica oleracea* L. var. *italica*) health-promoting compounds with inflorescence development. *Journal of Agricultural and Food Chemistry* **51(13)**: 3776-3782 (2003).

17. Kushad M, Brown AF, Kurilich AC, Juvik JA, Klein BP, Wallig MA, Jeffery EH, Variation of glucosinolates in vegetable crops of *Brassica oleracea*. *Journal of Agricultural and Food Chemistry* **47(4)**: 1541-1548 (1999).

18. Jones RB, Imsic M, Franz P, Hale G, Tomkins RB, High nitrogen during growth reduced glucoraphanin and flavonol content in broccoli (*Brassica oleracea* var. *italica*) heads. *Australian Journal of Experimental Agriculture* **47**: 1498-1505 (2007).

19. Cartea ME, Velasco P, Obregon S, Padilla G, de Haro A, Seasonal variation in glucosinolate content in *Brassica oleracea* crops grown in northwestern Spain. *Phytochemistry* **69**: 403-410 (2008).

20. Schonhof I, Klaring HP, Krumbein A, Claussen W, Schreiner M, Effect of temperature increase under low radiation conditions on phytochemicals and ascorbic acid in greenhouse grown broccoli. *Agriculture Ecosystems & Environment* **119(1-2)**: 103-111 (2007).

21. Jones RB, Effects of postharvest handling conditions and cooking on anthocyanin, lycopene, and glucosinolate content and bioavailability in fruits and vegetables. *New Zealand Journal of Crop and Horticultural Science* **35(2)**: 219-227 (2007).

22. Winkler S, Faragher J, Franz P, Imsic M, Jones R., Glucoraphanin and flavonoid levels remain stable during simulated transport and marketing of broccoli (*Brassica oleracea* var. *italica*) heads. *Postharvest Biology and Technology* **43(1)**: 89-94 (2007).

23. Lester GE, Manthey JA, Buslig BS, Organic vs. Conventionally Grown Rio Red Whole Grapefruit and Juice: Comparison of Production Inputs, Market Quality, Consumer

- 1
2
3 517 Acceptance, and Human Health-Bioactive Compounds. *Journal of Agricultural and Food*
4
5 518 *Chemistry* **55**: 4474-4480 (2007).
6
7 519 24. Soltoft M, Nielsen J, Laursen KH, Effects of Organic and Conventional Growth
8
9 520 Systems on the Content of Flavonoids in Onions and Phenolic Acids in Carrots and Potatoes.
10
11 521 *Journal of Agricultural and Food Chemistry* **58**: 10323–10329 (2010).
12
13 522 25. Lalor S, Coulter B, *Major and minor micronutrient advice for productive agricultural*
14
15 523 *crops*. 3rd ed. Teagasc, Oakpark, Carlow, 2008.
16
17 524 26. Alexander S, *Approved Pesticides for use on vegetable crops 2011*. Teagasc,
18
19 525 Kinsealy, Dublin 17, 2011.
20
21 526 27. Gardiner M.J, National vegetable and glasshouse crops research centre Kinsealy,
22
23 527 Malahide, Co. Dublin. *Soil Survey Bulletin* **3**: 1-8, 1963.
24
25 528 28. Reilly K, Cullen E, Lola-Luz T, Stone D, Valverde J, Gaffney, M., Brunton, N, Grant
26
27 529 J, Griffiths B, Effect of organic, conventional and mixed cultivation practices on soil
28
29 530 microbial community structure and nematode abundance in a cultivated onion crop.
30
31 531 *Journal of the Science of Food and Agriculture* **93(15)**: 3700-3709 (2013).
32
33 532 29. Marinova D, Ribarova F, Atanassova M, Total phenolics and total flavonoids in
34
35 533 Bulgarian fruits and vegetables. *J. Univ. Chem. Tech. Met.* **40(3)**: 255-260 (2005).
36
37 534 30. Hernandez-Hierro JM, Valverde J, Villacreces S, Reilly K, Gaffney M, Gonzalez-
38
39 535 Miret ML, Heredia FJ, Downey G, Feasibility Study on the Use of Visible-Near Infrared
40
41 536 Spectroscopy for the Screening of Individual and Total Glucosinolate Contents in Broccoli
42
43 537 *Journal of Agricultural and Food Chemistry* **60**: 7352-7358 (2012).
44
45 538 31. Naoumkina MA, Zhao QA, Gallego-Giraldo L, Dai XB, Zhao PX, Dixon RA,
46
47 539 Genome-wide analysis of phenylpropanoid defence pathways. *Molecular Plant Pathology*
48
49 540 **11(6)**: 829-846 (2010).
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

541 32. Brown AF, Yousef GG, Jeffrey EH, Klein BP, Wallig MA, Kushad M, Juvik JA,
542 Glucosinolate profiles in broccoli: Variation in levels and implications in breeding for cancer
543 chemoprotection. *Journal of the American Society for Horticultural Science* **127(5)**: 807-813
544 (2002).

545 33. Krumbein A, Schonhof I, Influence of temperature and irradiation on glucosinolates in
546 broccoli heads. *Biologically Active Phytochemicals in Food* **269**: 477-479 (2001).

547 34. Schreiner M, Vegetable crop management strategies to increase the quantity of
548 phytochemicals. *European Journal of Nutrition* **44 (2)**: 85-94 (2005).

549 35. Schonhof I, Krumbein A, Brückner B, Genotypic effects on glucosinolates and
550 sensory properties of broccoli and cauliflower. *Food* **48**: 25-33 (2004).

551 36. Reilly K, On-farm and fresh produce management, in *Handbook of plant food*
552 *phytochemicals: sources stability and extraction*, ed. by Tiwari, BK, Brunton N, Brennan C.
553 Wiley-Blackwell Publishing, pp. 201-225 (2012).

554 37. Zhang Y, Tang L, Discovery and development of sulforaphane as a cancer
555 chemopreventive phytochemical. *Acta Pharmacologica Sinica* **28(9)**: 1343-1354 (2007).

556 38. Marconett CN, Sundar SN, Tseng M, Tin AS, Tran KQ, Mahuron KM, Bjeldanes
557 LF, Firestone GL, Indole-3-carbinol down-regulation of telomerase gene expression requires
558 the inhibition of estrogen receptor-alpha and Sp1 transcription factor interactions within the
559 hTERT promoter and mediates the G(1) cell cycle arrest of human breast cancer cells.
560 *Carcinogenesis* **32(9)**: 1315-1323 (2011).

561 39 Khuda-Bukhsh AR, Das S, Saha SK, Molecular Approaches Toward Targeted
562 Cancer Prevention with Some Food Plants and Their Products: Inflammatory and Other
563 Signal Pathways. *Nutrition and Cancer* **66(2)**: 194-205 (2014).

564 40. Hoefkens C, Sioen I, Baert K, *et al.*, Consuming organic versus conventional
565 vegetables: The effect on nutrient and contaminant intakes. *Food and Chemical Toxicology*
566 **48(11)**: 3058-3066 (2010).

567 41. Hunter D, Foster M, McArthur JO *et al.*, Evaluation of the Micronutrient Composition
568 of Plant Foods Produced by Organic and Conventional Agricultural Methods. *Critical*
569 *Reviews in Food Science and Nutrition* **(51)6**: 571-582 (2011).

570

571

572

573

FIGURES

Figure 1. Total phenolic content (GAE mg/100g FW) in broccoli cv. 'Belstar' and cv. 'Fiesta' under different treatment combinations and in two harvest years (Panel A: year 1 and panel B: year 2). OS+OP means fully organic treatment and CS+CP means fully conventional treatment. Treatment codes: OS = organic soil treatment, CS = conventional soil treatment, OP = organic pest-control, CP = conventional pest-control. Bars show the mean and standard error of field replicates (n = 4).

Figure 2. Total flavonoid content (CE mg/100g FW) in broccoli cv. 'Belstar' and cv. 'Fiesta' under different treatment combinations and in two harvest years (Panel A: year 1 and panel B: year 2). OS+OP means fully organic treatment and CS+CP means fully conventional treatment. Treatment codes: OS = organic soil treatment, CS = conventional soil treatment, OP = organic pest-control, CP = conventional pest-control. Bars show the mean and standard error of field replicates ($n = 4$).

Figure 3. Representative electrochromatogram of desulfo-glucosinolates from broccoli. Each number above the peak represents a major identified desulfo-glucosinolate. **1-** desulfo-glucoraphanin; **2-**desulfo-protoitrin; **3-** desulfo-sinigrin; **4-** desulfo-glucotropaeolin (used as internal standard); **5-** desulfo-glucobrassicin; **6-** desulfo-4MeO-glucobrassicin; **7-** desulfo-neoglucobrassicin.

Figure 4. Average concentrations ($\mu\text{mol/g DW}$) of desulfo-glucobrassicin under different treatment combinations and two harvest years (Panel A: year 1 and panel B: year 2). OS+OP means fully organic treatment and CS+CP means fully conventional treatment. Treatment codes: OS = organic soil treatment, CS = conventional soil treatment, OP = organic pest-control, CP = conventional pest-control. Bars show the mean and standard error of field replicates ($n = 4$).

Figure 5. Average concentrations ($\mu\text{mol/g DW}$) of desulfo-neoglucobrassicin under different treatment combinations and two harvest years (Panel A: year 1 and panel B: year 2). OS+OP means fully organic treatment and CS+CP means fully conventional treatment. Treatment codes: OS = organic soil treatment, CS = conventional soil treatment, OP = organic pest-control, CP = conventional pest-control. Bars show the mean and standard error of field replicates (n = 4).

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TABLES

Table 1. Different agricultural management treatments applied in this study.

	Organic	Conventional
Soil treatment	<ul style="list-style-type: none">• Four year rotation: ley (red clover) → broccoli→ onion→ carrot• Additional organic fertilization as indicated by soil test• Winter cover crop	<ul style="list-style-type: none">• No set rotation, plots randomly allocated each year• Mineral fertilizers as indicated by soil test• No ley crop• No winter cover crop
Pest-control treatment	<ul style="list-style-type: none">• Certified organic seed• Refuge area• Mechanical pest-control• Weed control by mechanical methods• Certified organic treatments (e.g. garlic spray)	<ul style="list-style-type: none">• Chemically treated seed• Chemical weed control (herbicides)• chemical pest – control (fungicides and insecticides)

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Table 2. Specific pest-control and soil treatments used for broccoli cultivation in the Teagasc Kinsealy Systems Comparison trial in 2009 (year1) and 2010 (year 2).

	YEAR 1 (2009)	YEAR 2 (2010)
PEST-CONTROL TREATMENT		
<u>Organic Pest-control (OP)</u>	ECOGuard garlic spray modular drench ³ (4%v/v at 2L/m ²) Brassica collars Mechanical weeding (hand hoeing). Pyrethrum SEC ³ (1.1L/ha)	ECOGuard garlic spray modular drench ³ (4%v/v at 2L/m ²) Brassica collars Mechanical weeding (hand hoeing). Pyrethrum SEC ³ (1.1L/ha)
<u>Conventional Pest-control (CP)</u>	Dursban modular drench ³ (50ml per 5000 modules), Roundup ¹ (4L/ha), Stomp (3.3L/ha), Butisan S ¹ (1.5L/ha), Aramo ¹ (1.5L/ha), Decimate ¹ 20L/ha, Decis ³ (300ml/ha)	Proplant modular drench (2.4ml in 0.8L per 216 tray), Dursban modular drench ³ (50ml per 5000 modules), Roundup ¹ (4L/ha), Stomp (3.3L/ha), Butisan S ¹ (1.5L/ha), Stratos Ultra ¹ (3L/ha), Decis ³ (300ml/ha)
SOIL TREATMENT		
<u>Organic Soil treatment (OS)</u>	Previous crop – grass set aside >10 years N 115 kg/ha P 64 kg/ha K 180 kg/ha B 11kg/ha Applied as Greenvale plant food (4.5:3:3) (pelleted chicken manure + calcified seaweed), ProKali (3:0:14) and Solubor. A top dress equivalent to 20kg/ha N was applied on 14th August.	Previous crop – Red clover (2009) N 115 kg/ha P 64 kg/ha K 180 kg/ha B 11kg/ha Applied as Greenvale plant food (4.5:3:3) (pelleted chicken manure + calcified seaweed), ProKali (3:0:14) and Solubor.
<u>Conventional Soil treatment (CS)</u>	Previous crop – grass set aside >10 years N 115 kg/ha P 64 kg/ha K 180 kg/ha B 11kg/ha Applied as CAN (27% N), Single superphosphate (7.8%P), Sulphate of potash (42% K) and Solubor. A top dress equivalent to 20kg/ha N was applied on 14th August.	Previous crop – Any (2009) N 115 kg/ha P 64 kg/ha K 180 kg/ha B 11kg/ha Applied as CAN (27% N), Single superphosphate (7.8%P), Sulphate of potash (42% K) and Solubor.

¹ Herbicide, ² Fungicide, ³ Insecticide.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

6

For Peer Review

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3. Broccoli floret weight, total phenolic and total flavonoid content under different management practices.

	Mean floret weight (g)		Total phenolic content (GAE mg/100g FW)		Total flavonoid content (CE mg/100g FW)	
Treatment:	Year 1	Year 2	Year 1	Year 2	Year 1	Year 2
V1 + OS + OP	252.9±27.2	442.3±19.9	345.7±51.3	88.5±24.0	16.6±6.9	29.2±4.5
V1 + CS + OP	268.4±22.3	383.3±20.8	290.8±3.9	79.3±9.6	10.2±1.8	27.9±2.2
V1 + OS + CP	235.8±12.3	449.2±19.9	365.9±50.2	74.9±19.1	25.9±3.4	27.8±4.9
V1 + CS + CP	293.2±33.7	363.1±19.5	326.8±22.4	69.3±11.5	17.2±3.1	25.6±1.2
V2 + OS + OP	230.6±20.3	300.0±7.3	376.5±19.0	87.6±13.5	17.8±6.0	34.3±2.8
V2 + CS + OP	307.±17.0	247.1±78.5	273.4±16.0	106.8±7.5	5.6±1.1	35.4±5.2
V2 + OS + CP	253.7±8.5	327.2±24.3	375.6±63.0	70.2±19.1	22.0±7.3	23.8±3.5
V2 + CS + CP	293.9±41.4	363.9±15.2	267.0±18.4	86.7±15.4	11.9±2.1	29.8±2.0
Statistical significance:						
ANOVA p values:						
Replicate	0.9501	0.7573	0.9645	0.1610	0.4220	0.5511
variety	0.5924	0.0242	0.7736	0.3971	0.5049	0.4315
soil	0.0060	0.0775	0.0135	0.5927	0.0152	0.6748
pest-control	0.9044	0.3604	0.7419	0.2070	0.1627	0.1025
variety x soil	0.4692	0.1507	0.2991	0.2031	0.6107	0.2284
variety x pest-control	0.9785	0.0845	0.5692	0.7178	0.6720	0.1655
soil x pest-control	0.9329	0.4742	0.9263	0.9814	0.9793	0.6376
variety x soil x pest-control	0.2049	0.1905	0.8488	0.8701	0.7489	0.4929
fully conventional vs. fully organic	0.1698	0.8379	0.1600	0.4662	0.6037	0.1995

Data shown are mean ± standard error of the mean (n =4). OS+OP means fully organic treatment and CS+CP means fully conventional treatment. Since the difference between years was significant data for individual years is shown separately. Treatment codes: V1= cv. ‘Belstar’, V2 = cv. ‘Fiesta’ OS = organic soil treatment, CS = conventional soil treatment, OP = organic pest-control, CP = conventional pest-control. ANOVA p values in bold are significant at p<0.05.

Table 4. Broccoli total and individual glucosinolate content ($\mu\text{mol/g DW}$) under different management practices.

Treatment	Total glucosinolates	Glucoraphanin	Sinigrin	Glucobrassicin	4-MeOglucobrassicin	Neoglucobrassicin
V1 + OS + OP	9.26 $\pm$ 0.57	3.22 $\pm$ 0.21	0.37 $\pm$ 0.05	4.54 $\pm$ 0.21	0.36 $\pm$ 0.03	1.73 $\pm$ 0.13
V1 + CS + OP	10.09 $\pm$ 0.41	3.04 $\pm$ 0.22	0.38 $\pm$ 0.03	4.66 $\pm$ 0.29	0.35 $\pm$ 0.02	1.82 $\pm$ 0.16
V1 + OS + CP	9.01 $\pm$ 0.63	3.30 $\pm$ 0.25	0.40 $\pm$ 0.06	3.96 $\pm$ 0.36	0.33 $\pm$ 0.03	1.43 $\pm$ 0.18
V1 + CS + CP	9.06 $\pm$ 0.46	3.15 $\pm$ 0.28	0.40 $\pm$ 0.03	4.04 $\pm$ 0.30	0.35 $\pm$ 0.02	1.30 $\pm$ 0.13
V2 + OS + OP	9.99 $\pm$ 0.70	2.59 $\pm$ 0.26	0.94 $\pm$ 0.55	3.90 $\pm$ 0.39	0.26 $\pm$ 0.03	2.91 $\pm$ 0.28
V2 + CS + OP	9.15 $\pm$ 0.41	1.78 $\pm$ 0.19	0.65 $\pm$ 0.16	3.99 $\pm$ 0.24	0.25 $\pm$ 0.02	3.10 $\pm$ 0.21
V2 + OS + CP	8.38 $\pm$ 0.35	2.79 $\pm$ 0.32	0.50 $\pm$ 0.12	3.43 $\pm$ 0.20	0.19 $\pm$ 0.02	2.02 $\pm$ 0.18
V2 + CS + CP	8.09 $\pm$ 0.67	2.83 $\pm$ 0.34	1.64 $\pm$ 1.10	2.92 $\pm$ 0.23	0.19 $\pm$ 0.02	1.58 $\pm$ 0.16
Statistical significance:						
ANOVA p values:						
year	0.8272	0.0636	0.4388	0.8331	0.1054	0.2903
block	0.7676	0.0898	0.5193	0.6320	0.6335	0.7252
variety	0.2270	0.0329	0.0466	0.1872	0.0183	0.1223
soil	0.8776	0.2880	0.4852	0.5301	0.6759	0.2416
pest-control	0.2083	0.1182	0.6225	0.0670	0.3352	0.0197
variety x soil	0.2194	0.8252	0.5883	0.4062	0.5748	0.3861
variety x pest-control	0.4112	0.2318	0.5160	0.9482	0.3734	0.0145
soil x pest-control	0.9133	0.3195	0.0927	0.6916	0.3135	0.2475
variety x soil x pest-control	0.3401	0.3399	0.0501	0.5361	0.7351	0.6208
fully conventional vs. fully organic	0.1847	0.6651	0.3407	0.0187	0.2501	<.0001

Data shown are mean $\pm$ standard error of total and individual glucosinolates over two trial years ($\mu\text{mol/g DW}$). OS+OP means fully organic treatment and CS+ CP means fully conventional treatment. Treatment codes: Variety V1= 'Belstar', V2 = 'Fiesta' ; OS = organic soil treatment, CS = conventional soil treatment; OP = organic pest-control, CP = conventional pest-control. ANOVA p values shown in bold are significant at $p < 0.05$

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

For Peer Review

Supplementary material. Table S1. Climatic conditions during broccoli crop production in 2009 and 2010. T = Mean temperature (°C), TM = Mean maximum temperature (°C), Tm = Mean minimum temperature (°C), H = Mean humidity (%), V = Mean wind speed (Km/h), RA = Daily indicator for occurrence of rain or drizzle (total days), PP = Total monthly precipitation amount (mm).

	Month	T	TM	Tm	H	V	RA	PP
2009	April	8.8	12.4	4.3	84.0	17.9	22	70.1
	May	11.0	15.1	6.9	80.4	22.5	24	74.7
	June	13.8	18.2	8.4	78.6	16.6	19	65.0
	July	14.7	18.8	10.8	82.6	18.4	30	162.8
	August	15.4	19.3	11.4	81.4	20.7	28	69.3
	September	12.6	16.7	8.5	84.0	18.8	18	23.1
	Season	12.7	16.8	8.4	81.8	19.2	141	465.1
	(mean or total)							
	Month	T	TM	Tm	H	V	RA	PP
2010	March	5.2	9.7	0.1	81.1	17.2	17	55.6
	April	8.1	12.9	2.5	78.1	15.6	14	26.7
	May	10.2	14.7	4.8	78.3	15.3	20	29.2
	June	14.5	19.2	8.9	78.8	14.5	17	57.4
	July	15.7	19.6	12.1	80.4	19.5	28	76.7
	August	14.0	18.3	9.6	80.8	17	22	47.5
	Season	11.3	15.7	6.3	79.6	16.5	118	293.1
	(mean or total)							

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Year 1: Kinsealy Systems Comparison Trial

Supplementary material. Figure S1. Diagram showing the Kinsealy Systems Comparison Trial in Year 1. V1 = Variety 1, V2 = Variety 2 (for broccoli V1 = ‘Belstar’, V2 = ‘Fiesta’). CP = Conventional pest-control treatment, OP = Organic pest-control treatment. CS= Conventional soil treatment, OS= Organic soil treatment. Dark grey areas indicate 1 m wide grass inter-plot areas. Light grey indicates permanent refuge areas planted with buckwheat, vetch, borage, sunflower, coriander, fennel, cornflower, corn marigold, cocksfoot grass to encourage beneficial insects as part of the OP treatment. Note: Crops within the organic soil treatment (OS) move each year according to a set rotation (Ley → broccoli→onion→carrot). Crops within conventional soil treatment (CS) do not follow a set rotation pattern and move randomly, taking into account practices of commercial growers e.g. onion 4 year gap before returning to same land (1 in every 4). The “other” crop used in CS plots in year 1 was wheat (*Triticum aestivum*) and in year 2 was lettuce (*Lactuca sativa*).

Year 2: Kinsealy Systems Comparison Trial

Supplementary material. Figure S2. Diagram showing the Kinsealy Systems Comparison Trial in Year 2. V1 = Variety 1, V2 = Variety 2 (for broccoli V1 = 'Belstar', V2 = 'Fiesta'). CP = Conventional pest-control treatment, OP = Organic pest-control treatment. CS= Conventional soil treatment, OS= Organic soil treatment. Dark grey areas indicate 1 m wide grass inter-plot areas. Light grey indicates permanent refuge areas planted with buckwheat, vetch, borage, sunflower, coriander, fennel, cornflower, corn marigold, cocksfoot grass to encourage beneficial insects as part of the OP treatment. Note: Crops within the organic soil treatment (OS) move each year according to a set rotation (Ley → broccoli → onion → carrot). Crops within conventional soil treatment (CS) do not follow a set rotation pattern and move randomly, taking into account practices of commercial growers e.g. onion 4 year gap before returning to same land (1 in every 4). The "other" crop used in CS plots in year 1 was wheat (*Triticum aestivum*) and in year 2 was lettuce (*Lactuca sativa*).