

# Journal Pre-proof

Delivery of  $\beta$ -carotene to the *in vitro* intestinal barrier using nanoemulsions with lecithin or sodium caseinate as emulsifiers

Ariadna Gasa-Falcon, Elena Arranz, Isabel Odriozola-Serrano, Olga Martín-Belloso, Linda Giblin



PII: S0023-6438(20)31048-3

DOI: <https://doi.org/10.1016/j.lwt.2020.110059>

Reference: YFSTL 110059

To appear in: *LWT - Food Science and Technology*

Received Date: 19 March 2020

Revised Date: 7 August 2020

Accepted Date: 11 August 2020

Please cite this article as: Gasa-Falcon, A., Arranz, E., Odriozola-Serrano, I., Martín-Belloso, O., Giblin, L., Delivery of  $\beta$ -carotene to the *in vitro* intestinal barrier using nanoemulsions with lecithin or sodium caseinate as emulsifiers, *LWT - Food Science and Technology* (2020), doi: <https://doi.org/10.1016/j.lwt.2020.110059>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Ltd.

**Ariadna Gasa-Falcon:** Investigation, Roles/Writing - original draft; Writing - review & editing

**Elena Arranz:** Conceptualization, Investigation, Roles/Writing - original draft; Writing - review & editing, Supervision

**Isabel Odriozola-Serrano:** Conceptualization, Funding acquisition, Roles/Writing - original draft; Writing - review & editing, Supervision

**Olga Martín-Belloso:** Conceptualization, Funding acquisition, Roles/Writing - original draft; Writing - review & editing, Supervision

**Linda Giblin:** Conceptualization, Funding acquisition, Roles/Writing - original draft; Writing - review & editing, Supervision

Journal Pre-proof

1 **Delivery of  $\beta$ -carotene to the *in vitro* intestinal barrier using**  
2 **nanoemulsions with lecithin or sodium caseinate as emulsifiers**

3

4 Ariadna Gasà-Falcon<sup>1</sup>, Elena Arranz<sup>2</sup>, Isabel Odriozola-Serrano<sup>1</sup>, Olga Martín-Belloso<sup>1</sup>, Linda  
5 Giblin<sup>2\*</sup>

6 <sup>1</sup>Department of Food Technology, University of Lleida-Agrotecnio Centre, Lleida, Spain

7 <sup>2</sup>Department of Food Bioscience, Teagasc Food Research Centre, Moorepark, Fermoy, Co.  
8 Cork, Ireland

9

10 \*corresponding author:

11 E-mail: linda.giblin@teagasc.ie

12

13

14

15

16

17

18

19

20

21

22

23

## 24 **Abstract**

25 To increase the intestinal delivery of dietary  $\beta$ -carotene, there is a need to develop  
26 nanostructured food systems to encapsulate this fat soluble bioactive. The aim of this study was  
27 to evaluate the bioaccessibility and bioavailability across the intestinal barrier of  $\beta$ -carotene-  
28 enriched nanoemulsions stabilised with two emulsifiers (lecithin or sodium caseinate) by  
29 coupling an *in vitro* gastrointestinal digestion with two *in vitro* cell culture models (Caco-2 or co-  
30 culture of Caco-2/HT29-MTX). Nanoemulsions stabilised with lecithin had significantly higher  $\beta$ -  
31 carotene in the gastrointestinal digested micellar fraction, lower  $\beta$ -carotene in the Caco-2 (and  
32 Caco-2/HT29-MTX) apical compartment and significantly higher  $\beta$ -carotene in Caco-2 cellular  
33 content compared to  $\beta$ -carotene-enriched nanoemulsions stabilised with sodium caseinate.  
34 Finally, to assess anti-inflammatory activity of digested nanoemulsions, lipopolysaccharide  
35 stimulated macrophages were exposed to Caco- 2 basolateral samples with levels of TNF- $\alpha$   
36 and IL- $\beta$ , subsequently quantified. A TNF- $\alpha$  response from stimulated THP-1 macrophages was  
37 elicited by basolateral samples, regardless the emulsifier used to formulate nanoemulsions.  
38 This study demonstrated that  $\beta$ -carotene permeability is influenced by the food derived  
39 emulsifier used for stabilising nanoemulsions, indicating that composition may be a critical factor  
40 for  $\beta$ -carotene delivery.

41

42 Keywords:  $\beta$ -carotene, nanoemulsions, *in vitro* digestion, intestinal barrier

43

44

45

46

47

48

## 49 1. Introduction

50  $\beta$ -carotene is a vitamin A precursor with poor water-solubility (0.0006 g/L at 25°C). Positive  
51 health benefits associated with  $\beta$ -carotene consumption include lower incidence of cancer,  
52 cardiovascular diseases and degenerative disorders (Goralcyk, 2009). These health attributes  
53 have been related to  $\beta$ -carotene's antioxidant and immunomodulatory bioactivities, proven both  
54 *in vitro* (Bai et al., 2005) and *in vivo* (Zhou et al., 2018).

55 Although the recommended dietary allowance (RDA) for  $\beta$ -carotene has not been set, the U.S.  
56 Food and Drug Administration and the European Food Safety Authority derive a RDA for  
57 vitamin A of 900-700  $\mu$ g of retinol activity equivalents (RAE) daily and a population reference  
58 intake of 750-650  $\mu$ g REA daily, respectively. In addition,  $\beta$ -carotene appears to be degraded by  
59 the acidic environment of the stomach (Boon, McClements, Weiss, & Decker, 2010) which  
60 undoubtedly reduces  $\beta$ -carotene concentration in the intestine.

61 As a solution, recent studies have revealed the use of nanostructured delivery systems such as  
62 nanoemulsions to encapsulate and protect  $\beta$ -carotene after oral consumption and enhance its  
63 delivery to intestinal barrier (Chen, Li, Li, McClements, & Xiao, 2017; Gasa-Falcon et al., 2020;  
64 Yi, Zhong, Zhang, Yokoyama, & Zhao, 2015).  $\beta$ -carotene enriched nanoemulsions stabilised  
65 with pectin, lecithin, sodium caseinate, Tween 20 or sucrose palmitate, have been subjected to  
66 *in vitro* gastrointestinal digestion (GID) and the subsequent release of  $\beta$ -carotene has been  
67 determined (Gasa-Falcon et al., 2019; Salvia-Trujillo et al., 2013; Teixé-Roig et al., 2020).  
68 However, less information is currently available about the use of either proteins or  
69 phospholipids-based emulsifiers on  $\beta$ -carotene transit across the intestinal barrier and  
70 subsequent bioactivity from nanoemulsions post gastrointestinal digestion (GID). For instance,  
71 Lu, Kelly, & Miao (2017) described that the permeability of  $\beta$ -carotene in undifferentiated naïve  
72 gastrointestinal epithelial cells is depended on the emulsifier type added to the nanoemulsions  
73 rather than initial particle size of the nanoemulsions.

74 To model absorption *in vitro*, the use of differentiated monolayers expressing tight junctions,  
75 best represent the morpho-functional features of the intestinal barrier (Guri, Gülseren, &  
76 Corredig, 2013). However, *in vitro* monolayers present their own challenges since digestive  
77 fluids and nanoemulsions post GID (micellar fractions) are cytotoxic at relatively low

78 concentrations (Arranz, Corredig, & Guri, 2016). As a result, quantification of compounds on the  
79 basolateral side can be challenging.

80 Thus, the aim of this study was to evaluate the permeability of  $\beta$ -carotene-enriched  
81 nanoemulsions stabilised with two different emulsifiers across 21 days differentiated Caco-2  
82 and Caco-2/HT29-MTX co-cultures, post *in vitro* static GID. These 21 days old Caco-2/HT29-  
83 MTX co-cultures best represent the mature intestinal mucus barrier. Emulsifiers (lecithin and  
84 sodium caseinate) were selected based on their different properties (low (758 g/mol) and high  
85 ( $\approx$ 10-50 KDa) molecular weight), sources (synthetic and natural) and previous physiochemical  
86 characterisation of nanoemulsions stabilised with these emulsifiers in our group (Gasa-Falcon  
87 et al., 2019). Furthermore, the subsequent basolateral anti-inflammatory activity was assessed  
88 by quantification of TNF- $\alpha$  and IL- $\beta$  in lipopolysaccharide (LPS) stimulated macrophages (THP-1  
89 cells).

90

## 91 **2. Materials and methods**

### 92 **2.1. Materials**

93 Corn oil (Mazola, ACH Food Companies Inc., Memphis, TN) was purchased from a local  
94 market.  $\beta$ -carotene (synthetic,  $\geq$ 93% (UV), powder) was sourced from Sigma–Aldrich (Ireland).  
95 Lecithin was obtained from Alfa Aesar (Karlsruhe, Germany). Sodium caseinate (NaCas) ( $\geq$ 92%  
96 purity) was from Acros Organics (Geel, Belgium). The Caco-2 cell line was purchased from the  
97 European Collection of Cell Cultures (ECACC 86010202) and the human monocyte THP-1  
98 (ATCCTIB-202) and the human colon adenocarcinoma HT-29 cell lines (ATCCHTB-38) were  
99 purchased from American Type Culture Collection. This latter cell line was differentiated to HT-  
100 29-MTX following the protocol described by Guri et al. (2013). Tissue culture plastics were  
101 sourced from Sarstedt Ltd. (Wexford, Ireland). CellTiter 96 AQueus One Solution reagent was  
102 purchased from Promega (MyBio, Kilkenny, Ireland). Milli-Q water was used to prepare all  
103 nanoemulsions. All other chemicals were sourced from Sigma–Aldrich (Ireland) unless specified  
104 otherwise.

105

### 106 **2.2. Preparation of nanoemulsions**

107 Primary emulsions were prepared by mixing 4% (w/w) of the lipid phase (corn oil enriched with  
108 0.5% w/w of  $\beta$ -carotene) with 96% (w/w) of the aqueous phase containing the emulsifier (lecithin  
109 or NaCas) at 2% (w/w). Both phases were mixed with an Ultra-Turrax (IKA, Staufen, Germany)  
110 at 9500 rpm for 3 minutes. Then, primary emulsions were passed through an APV 1000 (SPX  
111 Flow Technology, Charlotte, NC, USA) at 500 bars for 3 cycles to obtain nanoemulsions.

112

### 113 **2.3. Determination of nanoemulsions** 114 **properties**

115 Particle size of nanoemulsions was determined using a Mastersizer 3000 (Malvern Instruments  
116 Ltd, Worcestershire, UK). The results were reported as the surface mean diameter ( $d_{43}$  ( $\mu\text{m}$ ))  
117 and the width of the distribution. The refractive index of the corn oil and water employed to  
118 perform the analysis were 1.34 and 1.33, respectively.

119 The emulsions  $\zeta$ -potential was determined using a Zetasizer NanoZS (Malvern Instruments Ltd,  
120 Worcestershire, UK). Samples were previously diluted (1/100) and equilibrated prior to  
121 analysis.

122 Physical stability of  $\beta$ -carotene enriched nanoemulsions was determined with an analytical  
123 centrifuge LUMiSizer 6112 (L.U.M. GmbH, Berlin, Germany) that accelerates destabilisation of  
124 samples. Results were analysed using the software package SEPView 6.0 (L.U.M. GMBH) that  
125 records transmitted light across the sample length and calculates the instability index that  
126 ranges from 0 to 1, with the greatest instability at 1. Instrumental parameters used for physical  
127 stability analysis were: speed 2,186 rcf; time interval 20 seconds; exposure time 10,000  
128 seconds; temperature 25°C.

129

### 130 **2.4. *In vitro* static simulated digestion**

131 Nanoemulsions were subjected to a simulated *in vitro* static GID (gastric and upper intestinal  
132 phases) that mimics the adult human upper gut. The INFOGEST standardised method  
133 (Brodkorb et al., 2019) was followed with minor modifications. Briefly, gastric phase consisted of  
134 5 mL of nanoemulsion with simulated gastric fluid containing porcine pepsin (EC 3.4.23.1)  
135 (3925.3 U/mg); pH was adjusted to 3.0 using HCl (1 M) and volume to 10 mL (Milli-Q water).  
136 The mixture was incubated for 2 h at 37°C with continuous shaking in a rotator. After 2h, pH

137 was increased to 6.5 using NaOH and 20  $\mu$ L  $\text{CaCl}_2$  (0.3 M), 4 ml bile (630 g/mol, EC232-369-0),  
138 and 2.5 mL pancreatin based on trypsin activity (8.13 U/mg; EC232.468.9) were added. Based  
139 on Verkempinck et al. (2017), extra lipase (pancreatin and lipase) was added to reach 420  
140 U/mL. The pH of the mixture was adjusted to 7.0, the volume to 20 mL with milli-Q water, and  
141 the mixture was incubated for 2 h at 37°C. The digestion was then stopped by adding protease  
142 inhibitor 4-(2-aminoethyl) benzenesulfonyl fluoride hydrochloride (1 mM). To obtain the micellar  
143 fraction, the digested fractions were centrifuged (Heraeus Megafuge 1.0, Massachusetts, USA)  
144 at 2890 x g for 40 minutes at 4 °C (Garrett, Failla, Sarama, & Craft, 1999). Samples were stored  
145 at -80°C for further experiments.

146

## 147 **2.5. *In vitro* cell based assays**

148 Caco-2 and HT29-MTX cell lines were grown in 75 cm<sup>2</sup> tissue culture flasks in a humidified 37  
149 °C incubator with a 5% CO<sub>2</sub> air atmosphere. Cells were cultured in Dulbecco's modified Eagle's  
150 medium (DMEM) supplemented with 10% (v/v) fetal bovine serum (FBS), 100 U/mL penicillin,  
151 and 100 mg/mL streptomycin. At 80% confluency, cells were trypsinated (0.25% trypsin/EDTA),  
152 diluted 1:6 in DMEM medium, and reseeded in flasks. Caco-2 and HT29-MTX cell lines in this  
153 study were used at passage number 29-41 and 53-67, respectively.

154

## 155 **2.6. Cytotoxicity of micellar fractions**

156 Caco-2 cells were seeded at a density of  $8 \times 10^4$  cells/well in 96-well plate. After 24 h of  
157 incubation, cells were washed with PBS. GID micellar fractions were filtrated (0.45  $\mu$ m), diluted  
158 in complete DMEM (between 2-16% v/v) and 80  $\mu$ L were added to each well. Subsequently, 20  
159  $\mu$ L of CellTiter 96<sup>®</sup> AQueous One Solution Cell Proliferation Assay reagent was added to each  
160 well and cells were incubated for 2 h. After 2 h, the quantity of formazan produced was  
161 measured spectrophotometrically at 490 nm in a microplate reader (Synergy HT BioTek,  
162 Winooski, VT, USA). Results were expressed as the percentage of cellular viability relative to a  
163 control group (cells with DMEM medium) versus the micellar fraction concentration (% v/v).  
164 Cytotoxicity of pure  $\beta$ -carotene dissolved in DMSO was also evaluated (0.05-10  $\mu$ g/mL).

165



## 2.7. Permeability across intestinal barriers

166 Caco-2 cells were seeded at a density of  $6 \times 10^4$  cells per insert in 12-well Transwell® plates  
167 (0.4  $\mu\text{m}$  pore size, 1.2 cm diameter, Costar, Cambridge, MA). In co-culture experiments, Caco-2  
168 and HT29-MTX were grown separately and then seeded at a ratio of 75:25, to a final density  $6 \times$   
169  $10^4$  cells per insert.

171 Culture media of each plate was changed every two days for 21 days. The integrity of the cell  
172 monolayer was monitored by measuring the transepithelial electrical resistance (TEER) ( $\Omega \cdot \text{cm}^2$ )  
173 using a Millicell-ERS Voltohmmeter (Merck Millipore, Carrigtwohill, County Cork, Ireland). On  
174 day 21, apical and basolateral compartments were washed three times with PBS and 470.6  $\mu\text{L}$   
175 and 1500  $\mu\text{L}$  DMEM were added to apical and basolateral compartments, respectively. Then,  
176 micellar fractions (29.4  $\mu\text{L}$ ) were added to apical compartment and incubated for 2 h. During this  
177 2 h, the TEER value did not change significantly (data not shown).

178 After permeability experiment, apical and basolateral samples were collected. Moreover, cell  
179 monolayer was washed three times with PBS, scraped and collected. Cells were centrifuged  
180 (Heraeus Megafuge 1.0) for 3 min at  $215 \times g$  and the supernatant discarded. Cells were stored  
181 at  $-80 \text{ }^\circ\text{C}$  for further analysis.

## 2.8. Determination of $\beta$ -carotene

184 Extraction of  $\beta$ -carotene from samples (apical, cells and basolateral) was performed as  
185 described by Yuan, Gao, Zhao, & Mao (2008) with minor modifications. Briefly, the samples  
186 were filtrated (0.45  $\mu\text{m}$ ) and mixed with ethanol and hexane, followed by a centrifugation (2890  
187  $\times g$ , 5 min,  $5^\circ\text{C}$ ) (Heraeus Megafuge 1.0). The upper fraction was collected, dried under  $\text{N}_2$  and  
188 stored at  $-80 \text{ }^\circ\text{C}$ . Each sample extract was dissolved in 200  $\mu\text{L}$  of the injection solvent  
189 acetonitrile (ACN): methanol (MeOH) 7:3 (v/v): acetone 6.7:3.3 (v/v) and filtered through 0.2  $\mu\text{m}$   
190 nylon filters (Millipore, Bedford, MA). High-performance liquid chromatography system (Waters  
191 Xevo TQ-S, Milford, USA) equipped with a photodiode array detector (HPLC-PDA) at 450 nm  
192 and a column ACQUITY UPLC® (C18 BEH 130 Å, 1.7  $\mu\text{m}$ , 2.1  $\times$  150 mm) (Waters) ( $30^\circ\text{C}$  and  
193 flow rate 0.85 mL/min) were used. . Mobile phase consisted of solvent A: ACN: MeOH 7:3 (v/v)  
194 and solvent B: water 100%, and the flow was isocratic (100% ACN/MeOH 7/3).  $\beta$ -carotene was

195 quantified by comparison with external standards. Results were reported as ng/mL and cellular  
196 uptake as percentage of detected  $\beta$ -carotene in cells versus apical samples at time 0 h.

197

## 198 **2.9. Anti-inflammatory activity of basolateral** 199 **samples**

200 Human monocytes THP-1 were cultured in RPMI 1640 culture medium supplemented with 10%  
201 (v/v) FBS, 100 U/mL penicillin, and 100 mg/mL streptomycin at 37 °C in a 5% CO<sub>2</sub> humidified  
202 atmosphere. Cells were added at a density of 5 x 10<sup>5</sup> cells/mL in 24 well plates. Differentiation  
203 to macrophages was induced by adding 1  $\mu$ g/ $\mu$ L 12-O-tetradecanoyl phorbol-13-acetate (TPA)  
204 to cells followed by 48 h incubation. After differentiation to macrophages, cells were washed  
205 with PBS. LPS (0.05  $\mu$ g/mL) and Caco-2 basolateral samples were added to each well. After 24  
206 h incubation, the culture medium was collected for subsequent TNF- $\alpha$  and IL-1 $\beta$  quantification.  
207 Release of TNF- $\alpha$  and IL-1 $\beta$  was measured in the supernatants of THP-1 cells using ELISA kits  
208 (R&D Systems, Minneapolis, USA), according to manufacturer's instructions. Multiscanner  
209 autoreader (Synergy HT BioTek) was used to read the absorbance of the plates at 450 nm.

210

## 211 **2.10. Statistical analysis**

212 Experiments were performed in triplicate on at least two different days and data was expressed  
213 as the mean with standard deviation. To determine the statistically significant differences  
214 between samples, one-way ANOVA followed by Bonferroni test ( $p \leq 0.05$ ) was conducted with  
215 SigmaPlot 11.0.

216

## 217 **3. Results and discussion**

### 218 **3.1. Physicochemical properties of nanoemulsions**

219 Both nanoemulsions containing lecithin and NaCas had a monomodal particle size distribution  
220 (Fig. 1A) and exhibited particle sizes in the nanometer range (0.35 and 0.29  $\mu$ m, respectively)  
221 (Table 1), in line with our previous published results (Gasa-Falcon et al., 2019). Nanoemulsions  
222 had negative  $\zeta$ -potential values, with lecithin displaying the highest negative value (Table 1).

223 NaCas-stabilised nanoemulsions exhibited the lowest end point instability index ( $0.603 \pm 0.006$ )  
224 compared with lecithin-stabilised nanoemulsions ( $0.773 \pm 0.001$ ) (Fig. 1B).  
225 Emulsifiers with a low molecular weight and/or with a high hydrophilic–lipophilic balance (HLB)  
226 are associated with a high efficiency at producing small particle sizes in oil-in-water emulsions  
227 (Jo & Kwon, 2014). Lecithin has an HLB value of 8 while NaCas has a value of 14. Thus, the  
228 intermediate-low HLB value of lecithin could explain why nanoemulsions stabilised with this  
229 emulsifier exhibited the highest particle size and instability index compared to nanoemulsions  
230 with NaCas (Iyer et al., 2015). In addition, the mass of NaCas ( $\approx 10$ -50 KDa) (Ozturk &  
231 McClements, 2016), its gelation behaviour (Rodriguez-Patino & Pilosof, 2011) and the thick  
232 interfacial layer covering oil droplets (McClements et al., 1993) undoubtedly contributed to its  
233 nanoemulsion stability over the accelerated centrifugation process. The large negative  $\zeta$ -  
234 potential values observed for lecithin and NaCas nanoemulsions (-58.81 mV and -53.41 mV,  
235 respectively) could be attributed to the phospholipid head groups from lecithin, and the fact that  
236 the nanoemulsion pH of  $\approx 6.5$  differs to the NaCas isoelectric point ( $pI=4.6$ ) respectively (Chang  
237 & McClements, 2016). Interestingly,  $\zeta$ -potential of lecithin stabilised  $\beta$ -carotene nanoemulsions  
238 became less negative after *in vitro* GID, while NaCas emulsions post GID have a stronger  
239 negative value (Gasa-Falcon et al., 2019). In that previous study, particle size after *in vitro* GID  
240 in lecithin nanoemulsions was higher compared to NaCas.

241

### 242 **3.2. Cytotoxicity of micellar fractions**

243 The maximum non-toxic concentration of micellar fractions was different depending on the  
244 emulsifier used (Fig. 2). Micellar fractions with lecithin showed no cell toxicity ( $>90\%$  cell  
245 viability) when Caco-2 cells were exposed to concentrations below 6% (v/v) (Fig. 2A), while for  
246 those containing NaCas no cell toxicity was observed at concentrations under 10% (v/v) (Fig.  
247 2B).

248 In agreement to our results, several studies have demonstrated that emulsifiers are cytotoxic in  
249 a concentration-dependent manner and that toxicity of nanoemulsions depends on the nature of  
250 emulsifier employed (Buyukozturk, Benneyan, & Carrier, 2010; Ujhelyi et al., 2012).  
251 Furthermore, Sadhukha, Layek & Prabha (2018) observed that the aqueous fraction of digested  
252 lipid-based delivery systems was responsible for cytotoxicity in MDCK kidney cells, reducing cell

253 viability by 40%. It has been previously reported that monoglycerides induce dose-dependent  
254 apoptosis in mammalian cells (murine thymocytes), which consisted on a rapid reduction in  
255 mitochondrial transmembrane potential, production of reactive oxygen species, among other  
256 processes (Philippoussos, Arguin, Fortin, Steff, & Hugo, 2002). In our study, micellar fractions of  
257 nanoemulsions were likely to contain lipid digestion products (i.e. free fatty acids and  
258 monoglycerides), and together with emulsifiers contributed to damage the cell integrity.

259 Similar cell viability results were obtained when control micellar fractions (without  $\beta$ -carotene)  
260 were tested in Caco-2 cells, but interestingly control micellar fractions derived from lecithin-  
261 stabilised nanoemulsions were 60% less toxic compared to micellar fractions with  $\beta$ -carotene.  
262 This suggests that  $\beta$ -carotene plays a role in cell cytotoxicity of Caco-2 monolayers. Indeed,  
263 Wooster et al. (2017) observed that the presence of  $\beta$ -carotene in LCT nanoemulsions ( $IC_{50}$ = 51  
264  $\mu$ g/mL) increased four times their toxicity in differentiated Caco-2 cells compared to empty  
265 nanoemulsions ( $IC_{50}$ = 257  $\mu$ g/mL). In contrast, a preliminary study by our laboratory indicated  
266 that pure  $\beta$ -carotene present in nanoemulsions was not cytotoxic to undifferentiated Caco-2  
267 cells (between 0.05  $\mu$ g/mL and 10  $\mu$ g/mL) (data not shown), suggesting that Caco-2 cell  
268 monolayers with tight junctions are more sensitive to GID  $\beta$ -carotene-enriched nanoemulsions  
269 than undifferentiated Caco-2 cells. Certainly, oxidation products of  $\beta$ -carotene could have been  
270 generated during *in vitro* GID, specifically due to the acidic pH of the gastric phase (Failla,  
271 Chitchumronchokchai, Ferruzzi, Goltz, & Campbell, 2014). Oxidation of  $\beta$ -carotene can produce  
272 carotenoid aldehyde breakdown products, which have documented toxic effects on numerous  
273 cell lines (K562, RPE 28 SV4 and ARPE-19) at concentrations between 10–20  $\mu$ M (Hurst, Saini,  
274 Jin, Awasthi, & Van Kuijk, 2005).

275 To investigate the bioavailability of  $\beta$ -carotene from nanoemulsions, permeability experiments  
276 were performed with lecithin and NaCas micellar fractions at a concentration of 6% (v/v).

277

### 278 **3.3. Permeability of $\beta$ -carotene nanoemulsions**

279 After *in vitro* GID,  $\beta$ -carotene concentration present in the micellar fractions was significantly  
280 higher in lecithin nanoemulsions compared to those prepared with NaCas (Table 2). There was  
281 a significant reduction of  $\beta$ -carotene in apical compartment after 2 h incubation compared to  
282 time zero regardless of emulsifier used. Interestingly after 2 h incubation, the apical of NaCas-

283 stabilised nanoemulsions had a significantly higher amount of  $\beta$ -carotene compared to lecithin-  
284 stabilised nanoemulsions.  $\beta$ -carotene content in cell lysates of Caco-2 cells was 3 times higher  
285 than in Caco-2/HT29-MTX co-cultures. In addition, the  $\beta$ -carotene concentration in Caco-2 cells  
286 lysates was significantly higher in lecithin-stabilised nanoemulsions (2.28%) compared with  
287 nanoemulsions containing NaCas (1.72%). Concentration of  $\beta$ -carotene was significantly lower  
288 in Caco-2/HT29-MTX co-cultures cell lysates ( $\geq 0.74\%$ ), with no significant differences between  
289 both emulsifiers.

290 The reason why NaCas nanoemulsions had less  $\beta$ -carotene in the micellar fraction, the apical  
291 sample at time zero and the cell lysate (Caco-2) may be explained by the fact that proteins such  
292 as NaCas, can interact hydrophobically with carotenoids and create complexes that act as a  
293 physical barrier for gastrointestinal digestive enzymes access (Wackerbarth, Stoll, Gebken,  
294 Pelters, & Bindrich, 2009). Hence,  $\beta$ -carotene may have remained entrapped within these  
295 complexes and not solubilised within mixed micelles, a process which is essential for  
296 permeability across the intestinal barrier (Baskaran, Sugawara, & Nagao, 2003). Moreover,  
297 Yang, Decker, Xiao, & McClements (2015) observed that the addition of 36 mg phospholipids  
298 (eg. 1,2-Dioleoyl-*sn*-glycero-3-phosphocholine) within the digestive fluids increased the degree  
299 of lipid digestion after *in vitro* GIT of vitamin E emulsions. This may explain why lecithin-  
300 nanoemulsions had higher concentrations of  $\beta$ -carotene within the micellar fraction. Another  
301 study observed that the maximum cellular uptake of carotenoids ( $\beta$ -carotene and lutein) in  
302 differentiated Caco-2 cells was obtained when micelles contained 50  $\mu\text{mol/L}$  of  
303 lysophosphatidylcholine (phospholipid derived from phosphatidylcholine present in lecithin)  
304 (Sugawara et al., 2001).

305 In agreement with the present study, Li, Arranz, Guri & Corredig (2017) reported a lower  
306 permeability of  $\beta$ -carotene from liposomes using 21-day old Caco-2/HT29-MTX co-cultures  
307 compared to Caco-2 monolayers. Interaction with mucus produced by HT29-MTX cell line  
308 reduces permeability of mucoadhesive lipophilic molecules, such as  $\beta$ -carotene (Sigurdsson,  
309 Kirch, & Lehr, 2013). Co-culturing Caco-2 cells with HT-29MTX adds a further layer of mucus  
310 complexity to more closely resemble the *in vivo* environment (Arranz, Corredig & Guri, 2016),  
311 but reduces permeability rates which may hamper compound detection. Thus, in our study, the

312 use of Caco-2/HT29-MTX cell line resulted in lower recovery of  $\beta$ -carotene in the cell lysates  
313 regardless of the emulsifier.

314 The failure to detect  $\beta$ -carotene in basolateral compartment underlined the limitations of the  
315 experiment due to upper concentration limits imposed by cytotoxicity data and inadequate  
316 sensitivity of detection instrumentation. Also,  $\beta$ -carotene may not have arrived at the basolateral  
317 within the 2 h incubation period.

318

### 319 **3.4. Immune functionality of basolateral samples**

320 Basolateral samples from permeability assays with  $\beta$ -carotene NaCas nanoemulsions and  $\beta$ -  
321 carotene lecithin nanoemulsions significantly increased TNF- $\alpha$  secretion (112% and 124%  
322 respectively,  $p < 0.05$ ) compared to basolateral sample control (positive control =  $2291 \pm 138$   
323 pg/mL of TNF- $\alpha$ ) from LPS activated THP-1 macrophages (Fig. 3A). However, IL-1 $\beta$  levels were  
324 unchanged regardless of basolateral samples ( $4769 \pm 145$  pg/mL) (Fig. 3B).

325 Previous studies have demonstrated that  $\beta$ -carotene reduces levels of TNF- $\alpha$  and IL-1 $\beta$  levels  
326 secreted from LPS-stimulated RAW264.7 cells (murine macrophage cell line) and from LPS-  
327 treated peritoneal macrophages (Li, Hong, & Zheng, 2019) as well as from serum of BALB/c  
328 mice, intraperitoneally injected with  $\beta$ -carotene (10 mg/kg) plus LPS (4 mg/kg) (Bai et al., 2005).  
329 This discrepancy with our results may be explained by the bypass of the gut and the use of  
330 different test material ( $\beta$ -carotene alone versus basolateral samples of Caco-2 monolayers  
331 treated with  $\beta$ -carotene-enriched nanoemulsions). Applying  $\beta$ -carotene directly to LPS  
332 stimulated THP-1 cells will dose dependently reduce secreted levels of TNF- $\alpha$  (data not shown).

333 It is important to note that our results do not confirm the presence of  $\beta$ -carotene or metabolites  
334 in the basolateral compartment. It is possible that other GID components in the micellar fraction  
335 may be capable of modulating TNF- $\alpha$ , although previous studies have shown that lecithin,  
336 sodium caseinate or emulsions with different fatty acid composition do not up-regulate cytokine  
337 production (Mukhopadhyaya et al., 2014, Reimund et al., 2004, Treede et al., 2009). However,  $\beta$ -  
338 carotene can be metabolised to high molecular weight products ( $\beta$ -apo-8'-carotenal,  $\beta$ -apo-10'-  
339 carotenal,  $\beta$ -apo-12'-carotenal,  $\beta$ -apo-14'-carotenal,  $\beta$ -apo-15'-carotenal) and short-chain  
340 products (hycyclocitral,  $\beta$ -ionone, ionene, 5,6-epoxy- $\beta$ -ionone, dihydroactinidiolide and 4-oxo-

341 ionone) (Siems et al., 2005), which themselves may directly or indirectly act as pro-inflammatory  
342 agents (Yeh, Wang, Chen, & Wu, 2009).

343

#### 344 **4. Conclusions**

345 Bioaccessibility of GID  $\beta$ -carotene in a Caco-2 model was enhanced when  $\beta$ -carotene-enriched  
346 nanoemulsions were stabilised with lecithin compared to those stabilised with NaCas. Caco-2  
347 basolateral samples from both nanoemulsions elicited a TNF- $\alpha$  response from stimulated THP-1  
348 macrophages. This study elucidates the importance of nanoemulsion composition for *in vitro*  
349 cellular permeability assays and the hurdles faced by concentration limits. Nanostructured food  
350 systems using lecithin as emulsifier might be a potential tool to increase uptake of dietary  $\beta$ -  
351 carotene.

352

353

#### 354 **Acknowledgements**

355 Financial support was received from the Fondo Europeo de Desarrollo Regional (FEDER) and  
356 Ministerio de Economía y Competitividad (project AGL2015-65975-R). EA receives funding from  
357 Enterprise Ireland (Career-FIT) and the European Union's Horizon 2020 research and  
358 innovation programme under the Marie Skłodowska-Curie grant agreement No 713654 (Project  
359 MF2018-0151 Food-BIBS). LG is funded by Science Foundation Ireland-16/RC/3835 VistaMilk.  
360 AGF thanks the Agència de Gestió d'Ajuts Universitaris I de Recerca (AGAUR) from the  
361 Catalan government (Spain) for the pre-doctoral grant.

362

#### 363 **References**

- 364 Arranz, E., Corredig, M., & Guri, A. (2016). Designing food delivery systems: Challenges related  
365 to the *in vitro* methods employed to determine the fate of bioactives in the gut. *Food and*  
366 *Function*, 7, 3319-36.
- 367 Bai, S.K., Lee, S.J., Na, H.J., Ha, K.S., Han, J.A., Lee, H., Kwon, Y.G., Chung, C.K., & Kim,  
368 Y.M. (2005).  $\beta$ -carotene inhibits inflammatory gene expression in lipopolysaccharide-stimulated

- 369 macrophages by suppressing redox-based NF- $\kappa$ B activation. *Experimental & Molecular*  
370 *Medicine*, 37, 323-334.
- 371 Baskaran, V., Sugawara, T., & Nagao, A. (2003). Phospholipids affect the intestinal absorption  
372 of carotenoids in mice. *Lipids*, 38(7), 705-711.
- 373 Boon, C. S., McClements, D.J., Weiss, J., & Decker E.A. (2010). Factors influencing the  
374 chemical stability of carotenoids in foods. *Critical Reviews in Food Science and Nutrition*, 50(6),  
375 515-32.
- 376 Brodkorb, A., Egger, L., Alminger, M., Alvito, P., Assunção R., Balance, S., Bohn, T., Bourlieu-  
377 Lacanal, C., Carrière, F., Clemente, A., Corredig, M., Dupont, D., Dufour, C., Edwards, C.,  
378 Golding, M., Karakaya, S., Kikhus, B., Le Feunteun, S., Lesmes, U., Macierzanka, A., Mackie,  
379 A.R., Martins, C., Marze, S., McClements, D.J., Ménard, O., Minekus, M., Portmann, R., Santos,  
380 C.N., Souchon, I., Singh, R.P., Vegarud, G.E., Wickham, M.S.J., Weitschies, W., & Recio, I.  
381 (2019). INFOGEST static *in vitro* simulation of gastrointestinal food digestion. *Nature Protocols*,  
382 14(4), 991-1014.
- 383 Buyukozturk, F., Benneyan, J.C., & Carrier, R.L. (2010). Impact of emulsion-based drug delivery  
384 systems on intestinal permeability and drug release kinetics. *Journal of Controlled Release*, 142  
385 (1), 22-30.
- 386 Chang, Y., & McClements, D.J. (2016). Influence of emulsifier type on the *in vitro* digestion of  
387 fish oil-in-water emulsions in the presence of an anionic marine polysaccharide (fucoidan):  
388 Caseinate, whey protein, lecithin or Tween 80. *Food Hydrocolloids*, 61, 92-101.
- 389 Chen, J., Li, F., Li, Z., McClements, D.J., & Xiao, H. (2017). Encapsulation of carotenoids in  
390 emulsion-based delivery systems: Enhancement of  $\beta$ -carotene water dispersibility and chemical  
391 stability. *Food Hydrocolloids*, 69, 49-55.
- 392 Failla, M.L., Chitchumronchokchai, C., Ferruzzi, M.G., Goltz, S.R., & Campbell, W.W. (2014).  
393 Unsaturated fatty acids promote bioaccessibility and basolateral secretion of carotenoids and  $\alpha$ -  
394 tocopherol by Caco-2 cells. *Food & Function*, 5(6), 1101-1112



- 395 Garrett, D.A., Failla, M.L., Sarama, R.J., & Craft, N. (1999). Accumulation and retention of  
396 micellar  $\beta$ -carotene and lutein by Caco-2 human intestinal cells. *Journal of Nutritional*  
397 *Biochemistry*, 10(10), 573-581.
- 398 Gasà-Falcon, A., Odriozola-Serrano, I., Oms-Oliu, G., Martín-Belloso, O. (2019). Impact of  
399 emulsifier nature and concentration on the stability of  $\beta$ -carotene enriched nanoemulsions  
400 during in vitro digestion. *Food & Function*, 10, 713-722.
- 401 Gasà-Falcon, A., Odriozola-Serrano, I., Oms-Oliu, G., Martín-Belloso, O. (2020).  
402 Nanostructured lipid-based delivery systems as a strategy to increase functionality of bioactive  
403 compounds. *Foods*, 9(3), 325.
- 404 Goralcyk, R. (2009).  $\beta$ -carotene, and lung cancer in smokers: review of hypotheses and status  
405 of research. *Nutrition and Cancer*, 61 (6), 767-774.
- 406 Guri, A., Gülseren, I., & Corredig, M. (2013). Utilization of solid lipid nanoparticles for enhanced  
407 delivery of curcumin in cocultures of HT29-MTX and Caco-2 cells. *Food & Function*, 4(9), 1410-  
408 1419.
- 409 Hurst, J.S., Saini, M.K., Jin, G.F., Awasthi, Y.C., & Van Kuijk, F.J.G.M. (2005). Toxicity of  
410 oxidized  $\beta$ -carotene to cultured human cells. *Experimental Eye Research*, 81(2), 239-243.
- 411 Iyer, V., Cayatte, C., Guzman, B., Schneider-Ohrum, K., Matuszak, R., Snell, A., Manohar-  
412 Rajani, G., McCarthy, M.P., & Muralidhara, B. (2015). *Human Vaccines and*  
413 *Immunotherapeutics*, 11(7), 1853-1864.
- 414 Jo, Y., & Kwon, Y.J. (2014). Characterization of  $\beta$ -carotene nanoemulsions prepared by  
415 microfluidization technique. *Food Science and Biotechnology*, 23(1), 107-113.
- 416 Li, R., Hong, P., & Zheng, X. (2019).  $\beta$ -carotene attenuates lipopolysaccharide-induced  
417 inflammation via inhibition of the NF- $\kappa$ B, JAK2/STAT3 and JNK/p38 MAPK signaling pathways  
418 in macrophages. *Animal Science Journal*, 90(1), 140-148.
- 419 Li, Y., Arranz, E., Guri, A., & Corredig, M. (2017). Mucus interactions with liposomes  
420 encapsulating bioactives: Interfacial tensiometry and cellular uptake on Caco-2 and co-cultures  
421 of Caco-2/HT29-MTX. *Food Research International*, 92, 128-137.

- 422 Lu, W., Kelly, A., & Miao, S. (2017). Bioaccessibility and cellular uptake of  $\beta$ -carotene  
423 encapsulated in model o/w emulsions: influence of initial droplet size and emulsifiers.  
424 *Nanomaterials*, 7(9), 282.
- 425 McClements, D.J., Dickinson, E., Dungan, S.R., Kinsella, J.E., Ma, J.G., & Povey, M. J. W.  
426 (1993). Effect of emulsifier type on the crystallization kinetics of oil-in-water emulsions  
427 containing a mixture of solid and liquid droplets. *Journal of Colloids and Interface Science*,  
428 160(2), 293-297.
- 429 Mukhopadhyaya, A., Noronha, N., Bahar, B., Ryan, M.T., Murray, B.A., Kelly, P.M., O'Loughlin,  
430 I.B., O'Doherty, J.V., & Sweeney, T. (2014). *Food Science and Nutrition*, 2(6), 712-723.
- 431 Ozturk, B., & McClements, D.J. (2016). Progress in natural emulsifiers for utilization in food  
432 emulsions. *Current Opinion in Food Science*, 7, 1-6.
- 433 Philippoussis, F., Arguin, C., Fortin, M., Steff, A.M., & Hugo, P. (2002). Cellular specificity  
434 related to monoglyceride-induced cell death. *Immunology Letters*, 83(3), 221-230.
- 435 Reimund, J.M., Scheer, O., Muller, C.D., Pinna, G., Duclos, B., & Baumann, R. (2004). *In vitro*  
436 modulation of inflammatory cytokine production by three lipid emulsions with different fatty acid  
437 compositions. *Clinical Nutrition*, 23(6), 1324-1332.
- 438 Rodriguez-Patino, J.M., & Pilosof, A.M.R. (2011). Protein-polysaccharide interactions at fluid  
439 interfaces. *Food Hydrocolloids*, 25(8), 1925-1937.
- 440 Sadhuka, T., Layek, B., & Prabha, S. (2018). Incorporation of lipolysis in monolayer permeability  
441 studies of lipid-based oral drug delivery systems. *Drug Delivery and Translational Research*,  
442 8(2), 375-386.
- 443 Salvia-Trujillo, L., Qian, C., Martín-Belloso, O., McClements, D.J. (2013). Influence of particle  
444 size on lipid digestion and  $\beta$ -carotene bioaccessibility in emulsions and nanoemulsions. *Food*  
445 *Chemistry*, 141(2), 1472-80.
- 446 Siems, W., Wiswedel, I., Salermo, C., Crifò, C., Augustin, W., Schild, L., Langhans, C.D., &  
447 Sommerburg, O. (2005).  $\beta$ -carotene breakdown products may impair mitochondrial functions-  
448 Potential side effects of high-dose  $\beta$ -carotene supplementation. *Journal of Nutritional*  
449 *Biochemistry*, 16(7), 358-397.

- 450 Sigurdsson, H.H., Kirch, J., & Lehr, C.M. (2013). Mucus as a barrier to lipophilic drugs.  
451 *International Journal of Pharmaceutics*, 453(1), 56-64.
- 452 Sugawara, T., Kushiro, M., Zhang, H., Nara, E., Ono, H., & Nagao, A. (2001).  
453 Lysophosphatidylcholine enhances carotenoid uptake from mixed micelles by Caco-2 human  
454 intestinal cells. *The Journal of Nutrition*, 131(11), 2921-2927.
- 455 Teixé-Roig, J., Oms-Oliu, G., Ballesté-Muñoz, S., Odriozola-Serrano, I., Martín-Belloso, O.  
456 Improving the *in vitro* bioaccessibility of  $\beta$ -carotene using pectin added nanoemulsions. *Foods*,  
457 9(4), 447.
- 458 Treede, I., Braun, A., Jeliaskova, P., Giese, T., Füllekrug, J., Griffiths, G., Stremmel, W., &  
459 Ehehalt, R. (2009). TNF-alpha-induced up regulation of pro-inflammatory cytokines is reduced  
460 by phosphatidylcholine in intestinal epithelial cells. *BMC Gastroenterology*, 13, 9-53.
- 461 Ujhelyi, Z., Fenyvesi, F., Váradi, J., Fehér, P., Kiss, T., Veszelka, S., Deli, M., Vecsernyés, M., &  
462 Bácskay, I. (2012). Evaluation of cytotoxicity of surfactants used in self-micro emulsifying drug  
463 delivery systems and their effects on paracellular transport in Caco-2 cell monolayer. *European*  
464 *Journal of Pharmaceutical Sciences*, 47(3), 564-573.
- 465 Verkempinck, S.H.E., Salvia-Trujillo, L., Moens, L.G., Charleer, L., Van Loey, A.M., Hendrickx,  
466 M.E., & Grauwet, T. (2017). Emulsion stability during gastrointestinal conditions effects lipid  
467 digestion kinetics. *Food Chemistry*, 246, 179-191.
- 468 Wackerbarth, H., Stoll, T., Gebken, S., Pelters, C., & Bindrich, U. (2009). Carotenoid-protein  
469 interaction as an approach for the formulation of functional food emulsions. *Food Research*  
470 *International*, 42(9), 1254-1258.
- 471 Wooster, T.J., Moore, S.C., Chen, W., Andrews, H., Addepalli, R., Seymour, R.B., & Osborne,  
472 S.A. (2017). Biological fate of food nanoemulsions and the nutrients they carry-internalisation,  
473 transport and cytotoxicity of edible nanoemulsions in Caco-2 intestinal cells. *RSC Advances*,  
474 7(64), 40053-40066.
- 475 Yang, Y., Decker, E.A., Xiao, H., & McClements, D.J. (2015). Enhancing vitamin E  
476 bioaccessibility factors impacting solubilization and hydrolysis of  $\alpha$ -tocopherol acetate  
477 encapsulated in emulsion-based delivery systems. *Food & Function*, 6(1), 84-97.

- 478 Yeh, S.L., Wang, H.M., Chen, P.Y., & Wu, T.C. (2009). Interactions of  $\beta$ -carotene and  
479 flavonoids on the secretion of pro-inflammatory mediators in an in vitro system. *Chemico-*  
480 *Biological Interactions*, 179(2-3), 386-393.
- 481 Yi, J., Zhong, F., Zhang, Y., Yokoyama, W., & Zhao, L. (2015). Effects of lipids on *in vitro*  
482 release and cellular uptake of  $\beta$ -carotene in nanoemulsion-based delivery systems. *Journal of*  
483 *Agricultural and Food Chemistry*, 63(50), 10831-10837.
- 484 Yuan, Y., Gao, Y., Zhao, J. & Mao, L. (2008). Characterization and stability evaluation of  $\beta$ -  
485 carotene nanoemulsions prepared by high pressure homogenization under various emulsifying  
486 conditions. *Food Research International*, 41(1), 61-68.
- 487 Zhou, L., Ouyang, L., Lin, S., Chen, S., Liu, Y.J., Zhou, W., & Wang, X. (2018). Protective role  
488 of  $\beta$ -carotene against oxidative stress and neuroinflammation in a rat model of spinal cord  
489 injury. *International Immunopharmacology*, 61, 92-99.

**Table 1.** Particle size ( $\mu\text{m}$ ) and  $\zeta$ -potential (mV) of nanoemulsions stabilised with different emulsifiers (LE: lecithin; NaCas: sodium caseinate). Differences among nanoemulsions were compared using one-way ANOVA followed by Bonferroni test. Different letters indicate statistically significant differences within the parameter tested ( $p < 0.05$ ).

**Table 2.**  $\beta$ -carotene concentration (ng/mL) quantified by high-performance liquid chromatography (HPLC-PDA) in micellar fractions (after *in vitro* digestion), apical samples and basolateral samples from permeability experiments after 2h incubation with micellar fractions obtained after *in vitro* digestion of nanoemulsions stabilised with lecithin (LE) and sodium caseinate (NaCas). ND = not detected, i.e., below detection limit of 10 ng/ml. Within a row, different lowercase letters indicate statistically significant differences ( $p < 0.05$ ) between emulsifiers. For an emulsifier, a statistical difference between apical  $t=0$  and apical  $t=2\text{h}$  is denoted by \*. Statistical analysis was performed using one-way ANOVA followed by Bonferroni test ( $*p < 0.05$ ).

**Fig 1.** (A) Particle size distribution and (B) instability profile of  $\beta$ -carotene-enriched nanoemulsions (0.02%  $\beta$ -carotene w/w, 4% corn oil w/w) stabilised with 2% of lecithin (LE) or NaCas: sodium caseinate (NaCas).

**Fig 2.** Cell viability (%) of Caco-2 cells after 2h incubation with micellar fractions (with  $\beta$ -carotene) (% v/v) and control micellar fractions (without  $\beta$ -carotene) (% v/v) obtained after *in vitro* digestion of nanoemulsions stabilised with different emulsifiers. Micellar fractions containing lecithin (A) and NaCas (B) were diluted with complete DMEM. Control cells (Ctrl) were grown in media with no treatment (100% viability). Different uppercase and lowercase letters indicate significant differences to control cells for micellar fractions and control micellar fractions, respectively. Statistical analysis was performed using one-way ANOVA followed by Bonferroni test ( $*p < 0.05$ ). Percentage of cell viability above 80% was considered as non-cytotoxic.

**Fig 3.** Effects of basolateral samples resulted from permeability experiments with  $\beta$ -carotene-enriched nanoemulsions emulsified with either 2% sodium caseinate (NaCas+) or lecithin (LE+) on the secretion ( $\% \pm \text{SEM}$ ) of TNF- $\alpha$  (A) and IL-1 $\beta$  (B) in lipopolysaccharide (LPS)-stimulated THP-1 cells. Positive controls (Ctrl +) were LPS-stimulated THP-1 cells and negative controls (Ctrl -) were non-stimulated THP-1 cells. Both controls were incubated with basolateral samples collected from control Caco-2 monolayers. Different letters indicate significant differences. Statistical analysis was performed using one-way ANOVA followed by Bonferroni test ( $*p < 0.05$ ).

1 **Table 1.**

<b>Emulsifier</b>	<b>Particle size (<math>\mu\text{m}</math>)</b>	<b><math>\zeta</math>-potential (mV)</b>
<b>LE</b>	$0.35 \pm 0.001^a$	$-58.81 \pm 2.56^a$
<b>NaCas</b>	$0.29 \pm 0.001^b$	$-53.41 \pm 1.83^b$

2

3



4



Journal Pre-proof



Table 2.

Emulsifier	Caco-2 model (ng/mL)		Caco-2/HT29-MTX model (ng/mL)	
	LE	NaCas	LE	NaCas
Micellar fraction	36430 ± 665 <sup>a</sup>	32858 ± 70 <sup>b</sup>	37370 ± 521 <sup>a</sup>	32759 ± 102 <sup>b</sup>
Apical t=0h	2171 ± 39 <sup>a*</sup>	1930 ± 4 <sup>b*</sup>	2198 ± 40 <sup>a*</sup>	1927 ± 6 <sup>b*</sup>
Apical t=2h	251 ± 14 <sup>a</sup>	339 ± 65 <sup>b</sup>	263 ± 36 <sup>a</sup>	365 ± 44 <sup>b</sup>
Cells	49.6 ± 5.4 <sup>a</sup>	33.2 ± 8.4 <sup>b</sup>	15.1 ± 5.7 <sup>c</sup>	18.1 ± 11.2 <sup>c</sup>
Basolateral	ND	ND	ND	ND









## Highlights

- Permeability of  $\beta$ -carotene was assessed using *in vitro* cell culture models.
- Bioaccessibility of  $\beta$ -carotene was enhanced with lecithin-stabilised nanoemulsions
- Nanoemulsions elicited a TNF- $\alpha$  response from stimulated THP-1 macrophages
- This study elucidates the hurdles faced by concentration limits

Journal Pre-proof

The authors declare that there are no conflicts of interest.

Journal Pre-proof