

Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>

Contents lists available at [SciVerse ScienceDirect](#)

The Veterinary Journal

journal homepage: www.elsevier.com/locate/tvjl

Review

Bioexclusion of diseases from dairy and beef farms: Risks of introducing infectious agents and risk reduction strategies

John F. Mee^{a,*}, Tim Geraghty^b, Ronan O'Neill^c, Simon J. More^{d,e}^a Animal and Grassland Research and Innovation Centre, Teagasc Moorepark, Fermoy, Co. Cork, Ireland^b Scottish Centre for Production Animal Health and Food Safety, University of Glasgow Veterinary School, Garscube Estate, Glasgow, United Kingdom^c Department of Agriculture, Fisheries and Food, Central Veterinary Laboratory, Backweston, Co. Kildare, Ireland^d Herd Health and Animal Husbandry, UCD School of Agriculture, Food Science and Veterinary Medicine, University College Dublin, Belfield, Dublin 4, Ireland^e Centre for Veterinary Epidemiology and Risk Analysis, UCD School of Agriculture, Food Science and Veterinary Medicine, University College Dublin, Belfield, Dublin 4, Ireland

ARTICLE INFO

Article history:

Accepted 2 July 2012

Keywords:

Cattle

Bioexclusion

Biosecurity

Disease transmission

Management

ABSTRACT

Infectious disease represents a major threat to the productivity and welfare of cattle herds throughout the world. The introduction of infectious agents into dairy and beef farms may be through direct transmission (purchased cattle, reintroduced resident cattle and contact with contiguous cattle) or indirect transmission (fomites, visitors, other species, and biological materials) and this article reviews the evidence supporting these transmission routes. In the absence of eradication programmes for many endemic infectious diseases, bioexclusion is the key management process for risk reduction. Various ameliorative bioexclusion strategies have been recommended and the evidence supporting these protocols is considered.

© 2012 Elsevier Ltd. All rights reserved.

Introduction

Biosecurity has been defined as a strategy of management practices to prevent the introduction of diseases and pathogens to an operation and to control spread within the operation (Wells, 2000). Biosecurity comprises two components, namely, bioexclusion and biocontainment. Bioexclusion relates to preventive measures (risk reduction strategies) designed to avoid the introduction of pathogenic infections (hazards), whereas biocontainment relates to measures to limit within-farm transmission of infectious hazards and onward spread to other farms.

The implementation of bioexclusion plans on dairy or beef farms is voluntary in almost all countries (an exception being larger dairy farms in Denmark; Kristensen and Jakobsen, 2011). Therefore, farmers need to be motivated to both change existing behaviours and to continue to implement effective practices to avoid biosecurity breakdowns (Truyers et al., 2011). However, currently there is a lack of consensus internationally in the published literature regarding bioexclusion protocols (Daly, 2011; Moore et al., 2008), their efficacy (van Winden et al., 2005; Faust et al., 2001) and their cost-effectiveness. These barriers may explain the slow adoption of such practices by many farmers (Gunn et al., 2008; Heffernan et al., 2008; Moore et al., 2008).

There has been no recent published review focusing specifically on the bioexclusion aspects of biosecurity on European cattle farms. This gap was recently identified by Animal Health Ireland (AHI) as a key task which needed addressing prior to the development of farmer guidelines on biosecurity. AHI is a not-for-profit, partnership-based organisation providing national leadership and coordination of non-regulatory animal health issues in Ireland (More et al., 2011).

The objective of the present article is to examine the existing scientific literature on bioexclusion in dairy and beef cattle enterprises and to summarise current best practice based on that literature. The review focuses on routes of transmission, the risks associated with introduction of infection and how these risks can be reduced through bioexclusion. An empirical hierarchy of risks has been established from the published literature, cognisant of the fact that the relative importance of each risk varies with the disease. Added animals (either purchased or hired, and reintroduced resident cattle) and contiguous cattle represent the highest biosecurity risks, in that order of risk ranking (Van Winden et al., 2005; Lindberg and Alenius, 1999).

Purchased (and hired) cattle

Overview

Purchase of cattle, where incoming stock remains in direct contact with the recipient herd for an extended period of time,

* Corresponding author. Tel.: +353 25 42387.

E-mail address: john.mee@teagasc.ie (J.F. Mee).

Table 1
Risks (odds ratios) and outcomes of introducing specific infections with purchased cattle.

Infection	Outcome	Odds ratio	Case-control comparison (all compared to no purchase)	Reference
Bovine herpes virus-1	Seropositivity	1.4	Purchase of more than one cow	Van Winden et al. (2005)
Bovine virus diarrhoea virus	Infection	1.8	Purchase of a pregnant cow	Valle et al. (1999)
Contagious mastitis pathogens	Infection	1.4–1.6	Purchase of heifers or cows	Agger et al. (1994)
<i>Escherichia coli</i> O157	Infection	1.9	Purchase of cattle within the last 2 years	Schouten et al. (2004)
<i>Leptospira</i> spp.	Seropositivity	1.5–1.6	Purchase of cattle	Oliveira et al. (2010)
<i>Mycobacterium avium</i> subspecies <i>paratuberculosis</i>	Seropositivity	2.1	Purchase of a large number ($\geq 25\%$) of cattle	Wells (2000)
<i>Mycobacterium bovis</i>	Herd breakdown	2.0	Purchase from a market	Ramírez-Villaescusa et al. (2010)
<i>Mycoplasma bovis</i>	Seropositivity	10.8	Purchase of adult cattle	Burnens et al. (1999)
<i>Salmonella</i> spp.	Infection	3.4	Purchase of cattle	Van Winden et al. (2005)

presents the highest risk for introducing infectious hazards. Farming practices, such as hiring a bull and returning it after the breeding season, significantly increase the risk of entry of venereally-transmitted infections such as bovine virus diarrhoea (BVD), infectious bovine rhinotracheitis (IBR) and leptospirosis (Bishop et al., 2010; Sibley, 2010; Bazeley, 2011). The asymmetry of information between vendor and purchaser about seller herd health status may result in cattle for sale, particularly in markets (Chi et al., 2002a) with a higher prevalence of disease than in the general cattle population. The concept of asymmetrical information, with the seller knowing more about the product than the buyer, is described in detail by Akerlof (1970) as 'Akerlof's lemons'. The published empirical evidence supporting cattle purchase as a route of transmission of infection is summarised below on a disease-by-disease basis (see Table 1).

Bluetongue virus

Purchase of pregnant heifers from bluetongue restriction zones resulted in the introduction of the bluetongue virus (BTV-8) during the midge-free season through transplacental and contact transmission into an expanding naïve dairy herd in Northern Ireland, a BTV-8-free region (Menziez et al., 2008). Subsequent Dutch work confirmed a role for vertical transmission in the epidemiology of BTV-8 in cattle (Santman-Berends et al., 2010).

Bovine herpes virus-1 (BHV-1)

While cattle purchase is an important risk factor for the introduction of BHV-1 (van Schaik et al., 2002), interactions with herd size (purchase risk only detected for small herds with ≤ 50 cattle; Boelaert et al., 2005) and herd type (strong association for dairy herds and a weak association for mixed herds; Van Wuijckhuise et al., 1998) have been shown. Purchase of asymptomatic cattle of uncertain health status is a particular risk for BHV-1 introduction (Hemmingsen and Bitsch, 1983).

Bovine respiratory syncytial virus

Seroconversion and severe outbreaks of respiratory disease in isolated dairy herds caused by bovine respiratory syncytial virus have been associated with the purchase of new animals (Elvander, 1996). Imported cattle were linked to the introduction and consequent outbreaks of BRSV in multiple Norwegian cattle herds in 1995 (Uttenthal et al., 1996).

Bovine viral diarrhoea virus (BVDv)

A well-documented way of introducing BVDv into a herd is purchase of transiently or persistently infected cattle (Valle et al.,

1999), or of a dam ('Trojan cow') carrying a persistently infected (PI) fetus (Lindberg and Houe, 2005). Larger herds are at particular risk of BVDv introduction as they are more likely to be open (Bishop et al., 2010). Specific pathogen-free status does not necessarily prevent introduction of BVDv (van Schaik et al., 2002).

Digital dermatitis

Digital dermatitis increases markedly in incidence in expanding dairy herds (Faust et al., 2001), especially after restocking (Holliman, 2003). Buying replacement heifers is a particularly risky strategy. Rodriguez-Lainz et al. (1999) showed that farms which purchased heifers had a significantly higher prevalence of digital dermatitis than farms which did not and that there was a positive association between the risk of digital dermatitis and the number of heifers purchased.

E. coli 0157

The presence of *E. coli* 0157 on dairy farms has been shown to be significantly associated with recent purchase of animals (Schouten et al., 2004) and with the presence of large numbers of purchased animals (Nielsen et al., 2002).

Leptospira spp.

Animal purchase has been identified as a significant risk factor for herd seropositivity to any *Leptospira* serovar (Oliveira et al., 2010) and specifically for *Leptospira interrogans* serovar *hardjo* (van Schaik et al., 2002). The appearance of leptospirosis as a clinical problem has recently been attributed to importation of cattle from high seroprevalence countries (Jones, 2011).

Mastitis

New herd infections caused by transmission of contagious mastitis pathogens (e.g. *Mycoplasma* spp., *Nocardia asteroides*, *Staphylococcus aureus*, *Streptococcus agalactiae*) between herds, both nationally and internationally (Holliman, 2003), has been associated with recent purchase of heifers or cows, for example, following restocking after a disease outbreak (Agger et al., 1994; Cook and Holliman, 2004; Olde Riekerink et al., 2010). In addition, high numbers of cattle introductions (≥ 8) have been associated with a higher incidence of subclinical mastitis and higher bulk milk somatic cell count (Bruower et al., 2010).

Mycobacterium avium subspecies *paratuberculosis*

Purchase of bulls or large numbers ($\geq 25\%$) of breeding females (Wells, 2000), particularly within the previous 5 years (Pillars et al.,

2009) has been strongly associated with higher seroprevalence (Chi et al., 2002b; van Winden et al., 2005; Weering et al., 2005) and clinical incidence (Norton et al., 2009) of Johne's disease in dairy herds. Restocking after herd depopulation and importation of animals have also been associated with Johne's disease (Richardson et al., 2009; Barrett et al., 2011).

Mycobacterium bovis

Herd breakdowns due to bovine tuberculosis (bTB) have been found to be strongly associated with cattle movements (Gilbert et al., 2005). Purchase of stock, in particular from markets, has also been significantly associated with the risk of bTB breakdown on farm, even in areas endemic with bTB (Ramírez-Villaescusa et al., 2010). Gopal et al. (2006) described purchased animals as the most likely source of bTB introduction in 30/31 outbreaks discussed. These new introductions are an important cause of persistence as herds are re-exposed and possibly re-infected each time cattle are purchased.

Mycoplasma bovis

The risk of *Mycoplasma bovis* culture-positive bulk milk (Passchyn et al., 2012) and seropositivity in adults (Burnens et al., 1999), as well as clinical disease (both lameness and respiratory disease) in weanlings (Byrne et al., 2001) and in adults, have each clearly been shown to be associated with purchase of cattle.

Mycotic dermatitis

Ringworm (mycotic dermatitis) has been reported as a particular problem in imported naïve heifers contracting infection following purchase (Holliman, 2003). Papini et al. (2009) reported that prevalence rates of *Trichophyton verrucosum* were also higher when the cattle present in the farm were of mixed origin, rather than when the whole stock had been purchased or born in the farm (92% vs. 88.2% and 85.7%, respectively).

Neospora caninum

Between-herd movement of *Neospora caninum*-infected cattle significantly enhances the spread of *N. caninum*, particularly into low seroprevalence herds after restocking following a disease outbreak (Woodbine et al., 2008). Furthermore, Holliman (2003) described several *N. caninum* abortion outbreaks which followed the purchase of pregnant cattle and Björkman et al. (1996) described a *Neospora* herd problem which originated with two infected cows which were among the founding cows of the herd.

Salmonella spp.

Farmers who purchase cattle (asymptomatic latent carriers or persistent excretor calves or adults) are significantly more likely to introduce *Salmonella* spp. into their dairy herds (van Winden et al., 2005; Bergevoet et al., 2009), particularly where the purchase is recent (odds ratio [OR] 2.6 for purchase within 14 days compared to no purchase) and from dealers (OR 3.9 compared to purchase from other farmers) (Evans, 1996).

Overall, purchasing cattle is a significant risk factor for many infectious diseases. In their analysis of disease risks, Ortiz-Pelaez and Pfeiffer (2008) identified having an open herd to be the most significant risk factors (OR 6.2 compared to a closed herd) associated with the diagnosis of one or more of the following: *Mycobacterium bovis* infection, BVD, fasciolosis, BHV-1 infection, Johne's disease, mastitis, neosporosis, pneumonia, rotavirus infection and salmonellosis. The stress of transport and mixing may be an

additional risk factor in disease recrudescence, e.g. IBR and transmission (Bishop et al., 2010; Holliman, 2003).

Risk reduction strategies

Maintaining a closed herd is the most important biosecurity measure (van Winden et al., 2005) as it eliminates infection risk from purchased cattle. However, even specific pathogen-free and closed herds experience disease breakdowns (van Schaik et al., 2002), reflecting the important role of other transmission routes. Furthermore, self-containment is not always practical where herds are expanding and/or breeding of replacements is impossible. Hence alternative risk reduction strategies (pre-movement, movement and post-movement) have been proposed.

Pre-movement risk reduction strategies

Minimise the number of cattle purchased and the number of source herds

Reducing the number of animals purchased will reduce the risk of introduction of infectious agents (Collins et al., 2006; Davison et al., 2006; Bazeley, 2009). Reducing the number of herds from which the animals are purchased will also reduce the risk of selecting an animal from a high disease prevalence herd. Purchases from markets or dealers present a very high biosecurity risk (Ramírez-Villaescusa et al., 2010; Kristensen and Jakobsen, 2011).

Purchase cattle which have not previously bred

Purchasing non-pregnant animals reduces the risk of transmission of infection from an infected fetus, such as can occur with BVDv (Lindberg and Houe, 2005). Venereal diseases are more likely to be spread by animals which have bred; Sanderson et al. (2000) reported that using non-virgin bulls were associated with an increased risk of both trichomonosis and campylobacteriosis.

Purchase from herds with likely low disease prevalence

The disease prevalence in the source population of purchased cattle will greatly influence the risk of introduction of infectious agents. This principle may be applied at a national level, where farmers should avoid importing cattle from exotic disease-affected regions as well as at the farm level. Herds which are likely to have lower likely disease prevalence include: (1) Certified disease-free herds; the risk of introducing specific diseases can be greatly reduced by purchasing stock from certified disease-free source herds, (Wells, 2000; van Winden et al., 2005; Sibley, 2010) or high health status markets. (2) Closed herds; one key factor associated with increased disease prevalence is having an open rather than a closed herd (Chi et al., 2002b; Weering et al., 2005; Ortiz-Pelaez and Pfeiffer, 2008). (3) Smaller herds; for many diseases seroprevalence in cattle has been found to be proportionate to the size of the source population (Weering et al., 2005; Ortiz-Pelaez and Pfeiffer, 2008). One important example is bovine tuberculosis (bTB). In countries such as Ireland where bTB is endemic, herd risk is significantly associated with herd size (Olea-Poppelka et al., 2004; Wolfe et al., 2010).

Obtain cattle disease history

This may include a vendor or veterinary declaration of clinical disease incidence, medication and vaccine usage and previous laboratory results from the source herd (Bergevoet et al., 2009; Moore et al., 2009). Bergevoet et al. (2009) concluded that dairy herds could economically prevent the introduction of *Salmonella* spp. by excluding animals based on bulk milk antibody results. For diseases with long incubation periods (e.g. Johne's disease), pro-active assurances of freedom from clinical disease for 3 years or more is recommended (Pritchard, 1996).

Clinically examine the cattle for sale

While a clinical examination by the purchaser or their veterinary practitioner will not detect asymptomatic carrier animals, and the sensitivity of such examinations is variable, it is recommended for detection of the clinically obvious infectious disease (Pritchard, 1996; Bazeley, 2009; Moore et al., 2009). The examination may include cow-side examinations such as rectal temperature, body condition score, a California mastitis test (CMT), and udder, gait and reproductive examinations. However, such examinations applied singly have low sensitivity (Barkema et al., 2009), though for some diseases (for example, digital dermatitis) clinical examination is the only available diagnostic modality.

Quarantine cattle before movement

Quarantine has been defined as the isolation of cattle in an area that prevents direct contact with other livestock or the possibility of cross-contamination of animal waste (Davison et al., 2003). The recommended period of quarantine ranges from 3 (Maunsell and Donovan, 2008) to 6 weeks (Caldow, 2009), but commonly 4 weeks is advised (Bazeley, 2009) to cover the period of greatest risk of clinical disease after purchase, e.g. salmonellosis (Evans, 1996). Quarantine prior to sampling will improve the value of pre-purchase test results by allowing detectable seroconversion to an exposure that occurred immediately prior to quarantine. In addition, quarantine reduces the risk of post sampling infection on the farm of the vendor, and facilitates multiple observations and examinations.

Test cattle before movement

Laboratory testing prior to animal introduction is commonly recommended for many infectious diseases and can greatly enhance the sensitivity of detecting an infectious animal and therefore reduce risk (Wells, 2000; Maunsell and Donovan, 2008; Moore et al., 2009; Sibley, 2010). In general, the consequences of a false negative far outweigh those of a false positive; sensitivity should be favoured over specificity for pre-introduction tests (Barkema et al., 2009). Test insensitivity, especially in young animals, is a problem with Johne's disease detection but also in non-clinical adults where the probability of non-detection increases with the number of cattle purchased and greater prevalence (Carpenter et al., 2004; Collins et al., 2006). Pre-movement testing is not equally successful for all diseases, being proportionally less successful in BVD, IBR, Johne's disease and salmonellosis, in descending order (van Winden et al., 2005).

Medicate cattle before movement

In some cases, targeted medication can eliminate carrier status or reduce the probability of a carrier becoming infective. In such circumstances, incoming stock can be treated prophylactically to reduce the risk of introduction of infectious agents (preconditioning; Hilton and Olynk, 2011). Treatment with parenteral antibiotics, anthelmintic and flukicide, antibiotic foot-bathing and vaccination has been recommended (Bazeley, 2009), but may have a minimal effect depending on the disease (van Winden et al., 2005). Vaccination courses should be completed at least 2 weeks before release from quarantine (Caldow, 2009). Vaccination of resident cattle at the same time is also important as this can significantly reduce the risk of disease transmission (Nickell et al., 2011).

These strategies are discussed in the order which they are applied. However, if we were to rank the strategies in order of their likely impact our order would be: (1) information on the disease and biosecurity status of the source herd; (2) the number and history of the individuals purchased; (3) pre-introduction testing; (4) quarantine; and (5) medication.

Movement risk reduction strategies

The movement process itself may be targeted to reduce the risk of infection transmission by (1) loading and unloading animals at the perimeter of the farms; (2) not mixing animals from different sources in the transport vehicle; (3) transporting purchased cattle in the purchaser's vehicle (Maunsell and Donovan, 2008; Bazeley, 2009); and (4) minimising transport distances (Greger, 2007) and other stressors such as overcrowding. Legislation to control movement may also reduce the spread of infectious agents (Velthuis and Mourits, 2007), and is a frequent state-mediated reaction to exotic disease outbreaks.

Post-movement risk reduction strategies

Quarantine, testing and medication have been discussed as pre-movement strategies, but all may also be applied at the recipient farm (post-movement). This may reduce infection spread from stressed, clinically affected animals breaking down within weeks of movement, e.g. IBR, salmonellosis, Johne's disease, *Mycoplasma bovis* (Byrne et al., 2001; Bazeley, 2009). It is particularly important to prevent contact with breeding animals, e.g. BVDv (Sibley, 2010). Pregnant cattle should be group-segregated and isolated until calved (O'Farrell et al., 2001). Testing of the progeny of females purchased while pregnant can detect infectious agents transmitted transplacentally, e.g. BVDv and *Neospora* (Sibley, 2010).

Reintroduced resident cattle

Overview

Reintroduction of cattle returning from out-farms, common pastures, common accommodation, contract rearer accommodation, agricultural shows and marts or sales presents a risk of introducing infection. As they are not new additions to the herd, these animals are often not viewed as a threat to the resident herd disease status. However, in addition to the risk of comingling with infectious animals, there is a risk from contact with the external environment, handlers and the transport vehicle.

Co-grazing with cattle from other herds is a significant risk factor for introduction to the resident herd of BHV-1, BVDv (OR 3.4 compared to not sharing pasture), *Leptospira hardjo* (OR 1.63 compared to not sharing pasture) and *Salmonella* spp. (OR 8.9 compared to no co-mingling during contract heifer rearing) (Valle et al., 1999; van Schaik et al., 2002; Davison et al., 2006; Adhikari et al., 2009; Oliveira et al., 2010). Co-grazing with sheep and goats constitutes a risk for BVD, Johne's disease and leptospirosis (Caldow, 2004).

Sharing housing where cattle from different herds co-mingle is a significant risk factor for BVD (OR 15.1 compared to not sharing cattle housing) (Valle et al., 1999). Return of cattle following unsuccessful sale has been reported as a significant risk factor for introduction of BHV-1, BVDv, *E. coli* 0157, *Leptospira hardjo* and *Salmonella* Dublin to the resident herd (Cernicchiaro et al., 2009; van Schaik et al., 2002), although this may not always be the case: see, for example, Davison et al. (2006) who found no such relationship for *Salmonella* spp.). Participation in agricultural shows is a significant risk factor for introduction of BHV-1 (van Schaik et al., 2001), *Salmonella* spp. (Davison et al., 2003) and malignant catarrhal fever (MCF) (Moore et al., 2010).

Risk reduction strategies

Ameliorative strategies to reduce the risks associated with reintroduced cattle are in principle similar to those for purchased cattle. Key practices include (1) not using common pastures or shared

cattle housing; (2) not directly returning unsold cattle; (3) selling at high health status physical sales; (4) maintaining cattle-proof boundaries on out-farms; and (5) not participating in agricultural shows. Testing cattle prior to admittance to shows will lower, but does not eliminate, the risk of BHV-1 transmission (Breidenbach et al., 2005).

Contiguous contacts

Overview

Cattle are social animals with individuals and groups interacting whenever possible. Most farm boundaries have evolved to demarcate ownership of property and not for biosecurity purposes. The transmission significance of peripheral farm boundaries has been discussed regarding BHV-1 (van Schaik et al., 2001), bTB (O'Corry Crowe et al., 1996) and BVDv (van Schaik et al., 2002; Scharko et al., 2011). In addition, 71% of brucellosis outbreaks in Northern Ireland were most likely caused by direct contact between contiguous cattle at pasture (Abernethy et al., 2011).

Risk reduction strategies

To reduce the risks associated with contiguous contacts, the following strategies have been recommended based on the principle that there must be no opportunity for straying or for direct contact (Duncan, 1990): (1) Double spaced boundary fencing with a gap of 3 m has been proposed to reduce significantly the risk of spread of disease such as BVD or IBR (Crawshaw et al., 2002). There are no empirical recommendations published on fence height or design. (2) Electric scare wires on each side of the boundary fence or a strip of uninterrupted hedge or trees also ameliorate the risk of transmission of infection (Caldow, 2004); a higher density of hedgerows was found to be negatively correlated with bTB transmission (Mathews et al., 2006). (3) Vaccination against specific pathogens such as BVDv (Scharko et al., 2011).

In a study of contact between cattle farms in the UK, Brennan et al. (2008) found that over half of boundary fences perceived to prevent contact, nose-to-nose contact was in fact possible with animals on adjacent farms.

Fomites

Overview

Fomites have been implicated in the transmission of various cattle pathogens such as BVDv (Stevens et al., 2011), foot-and-mouth disease virus (Sutmoller et al., 2003), cryptosporidium (Nydam and Mohammed, 2005) and ovine herpesvirus 2 (the cause of MCF) (Moore et al., 2010). Contaminated farm machinery, visitor vehicles and veterinary equipment, (e.g. needles and syringes, nose tongs, halters, obstetrical equipment, dosing equipment, dehorning equipment, hoof paring equipment) can all act as mechanical vectors for introduction of infectious agents into herds (DiGiacomo et al., 1985; Gunn, 1993; Lang-Ree et al., 1994; Niskanen and Lindberg, 2003; Makoschey and Beer, 2004).

Risk reduction strategies

In order to reduce the risks posed by such fomites, farmers should use their own cattle equipment rather than veterinarians' equipment (Raaperi et al., 2010) and thoroughly clean, disinfect and rinse equipment before and after use (Morley, 2002).

Visitors

Overview

Visitors to farms are potential infection vectors particularly via their hands (Larson, 1995), but also their clothing, boots, equipment and vehicles (Morley, 2002; Kirk et al., 2003; Ellis-Iversen et al., 2011). Visitors can be categorised as low or high risk. High-risk visitors include collectors of dead stock, other farmers, veterinary practitioners, artificial insemination (AI) technicians, lay scanners and lay foot trimmers.

However, although visitors are known to be potential disease vectors, there is only limited published evidence which quantifies the risk of disease introduction from farm visitors. In a small study with a limited data-set, farms that had professional visitors who did not always use protective clothing had an increased risk of introducing BoHV-1 (OR 2.8) (van Schaik et al., 2001). Not providing boots for visitors has been shown to be a significant risk factor for seropositivity to bovine coronavirus and BRSV (Ohlson et al., 2010).

Risk reduction strategies

The key procedures to reduce the risks posed by high risk visitors are: (1) providing personal protective equipment, e.g. gloves, footwear and overalls (Morley, 2002; van Schaik et al., 2002); (2) restricting visitor contact only to the necessary stock; (3) installing a vehicle bath with appropriate disinfectant at the single farm entrance; (4) providing hand-washing and boot-washing facilities (disinfectant footmats have been shown to significantly reduce bacterial contamination of footwear; Amass, 2006; Dunowska et al., 2006); and (5) moving fallen stock to an area separated from livestock and farm activities and restricting collection service staff access only to this area.

Other species

Overview

Resident cattle may have indirect contact with other non-bovine domestic animals, feral animals, wildlife, vermin and humans (walkers, passing motorists, hunters, etc.). Wildlife is recognised as an important reservoir of infectious agents (Daszak et al., 2000; Simpson, 2002). For example, wild ruminants (deer and mouflon) act as reservoirs and may be involved in the dissemination and persistence of BTv (García-Bocanegra et al., 2011). Many wildlife species have been identified as important reservoirs for *Mycobacterium bovis*, including Eurasian badgers (*Meles meles*) in Ireland and the UK (More, 2009), white-tailed deer (*Odocoileus virginianus*) in the USA (Palmer et al., 2000) and the brushtail possum (*Trichosurus vulpecula*) in New Zealand (Jackson et al., 1995). *Mycobacterium avium* subsp. *paratuberculosis* (MAP) also has a wide host range, though the threat of spread to cattle from other species remains poorly understood (Greig et al., 1999; Beard et al., 2001).

Neospora caninum is a coccidian parasite of domestic dogs (and other canids) (Williams et al., 2000; King et al., 2011). *Cryptosporidium parvum* infection is prevalent in a range of mammalian and avian wildlife, resulting in environmental contamination and the potential for transmission to farmed livestock (Sturdee et al., 1999; Majewska et al., 2009; Samra et al., 2011). Wild deer are a potential source of infection for livestock of a broad range of bacteria, viruses and parasites (Böhm et al., 2007). *Clostridium botulinum* intoxication of cattle has been associated with poultry litter contamination of pasture (McLoughlin et al., 1998).

Risk reduction strategies

Several broad principles are relevant when seeking to reduce risk associated with other species, including providing and maintaining boundaries between known areas of significant wildlife populations and the other areas of the farm (these are not always the peripheral boundaries) and securely storing animal foodstuffs and animal wastes to remove potential attractants to wildlife (including birds). The measures necessary will vary depending on the geographic area and the wildlife species involved; for example, Ward et al. (2010) reviewed options to reduce the risk of *M. bovis* transmission from badgers to cattle.

Biological material

Overview

Introduction of biological material (colostrum, colostrum replacers, whole milk, semen, embryos, vaccines, slurry) into a farm is a potential biosecurity risk. For example, MAP can be transmitted in colostrum or whole milk, as can enzootic bovine leukosis (EBL) virus (Hopkins and DiGiacomo, 1997). Semen and embryos can transmit various infectious agents including BVDv and BHV-1, and cases of vaccine contamination with BVDv have been reported (Lindberg and Alenius, 1999; Barkema et al., 2001). Slurry and dirty water can be a biohazard with *E. coli* O157, *Salmonella* and *Campylobacter* surviving for up to 3 months (Nicholson et al., 2005). Despite being an obligate pathogen, MAP can survive in the environment for over 100 weeks (Whittington et al., 2004).

Risk reduction strategies

Avoiding introduction of biological material of uncertain health status is the best method of avoiding risk. Purchase of semen from AI centres approved for intercommunity trade and of embryos processed by internationally approved sanitary protocols will reduce disease introduction risks (EFSA, 2006; Givens and Marley, 2008). Pasteurisation of colostrum can reduce the load of infectious agents (Godden et al., 2006).

Conclusions

Bioexclusion is fundamental to the protection of herd health. This review has highlighted the relative importance of direct transmission of infectious agents via purchased cattle as the primary biosecurity risk and detailed the key protocols which can reduce this risk. Such protocols need to be prioritised and made farm-specific (Daly, 2011) based on the availability of resources (financial, labour, farm infrastructure) as well as the identified risks and consequences of infection introduction.

Conflict of interest statement

None of the authors of this paper has a financial or personal relationship with other people or organisations that could inappropriately influence or bias the content of the paper.

Acknowledgements

The authors thank the other members of the Animal Health Ireland Technical Working Group on Biosecurity who contributed to this document: Bosco Cowley (Schering Plough Intervet, Ireland), Bernard Eivers (National Cattle Breeding Centre, Enfield, Co. Meath, Ireland), Pat Kirwan (Unit 3C, Riverview Business Park, New Nangor Road, Dublin 12), Luke O'Grady (Herd Health and Animal

Husbandry, UCD School of Agriculture, Food Science and Veterinary Medicine, University College Dublin, Belfield, Dublin 4, Ireland), Regina Sayers (Animal and Grassland Research and Innovation Centre, Teagasc Moorepark, Fermoy, Co. Cork, Ireland), Michael Sexton (Riverview Veterinary Clinic, Bandon, Co. Cork, Ireland).

References

- Abernethy, D.A., Moscard-Costello, J., Dickson, E., Harwood, R., Burns, K., McKillop, E., McDowell, S., Pfeiffer, D.U., 2011. Epidemiology and management of a bovine brucellosis cluster in Northern Ireland. *Preventive Veterinary Medicine* 98, 223–229.
- Adhikari, B., Besser, T.E., Gay, J.M., Fox, L.K., Davis, M.A., Cobbold, R.N., Berge, A.C.B., Hancock, D.D., 2009. The role of animal movement, including off-farm rearing of heifers, in the interherd transmission of multidrug-resistant *Salmonella*. *Journal of Dairy Science* 92, 4229–4238.
- Agger, J.F., Priou, C., Huda, A., Aagaard, K., 1994. Risk factors for transmission of *Streptococcus agalactiae* infection between Danish dairy herds: A case control study. *Veterinary Research* 25, 227–234.
- Akerlof, G.A., 1970. The market for 'lemons': Quality uncertainty and the market mechanism. *Quarterly Journal of Economics* 84, 488–500.
- Amass, S.F., 2006. Effectiveness of using a mat filled with a peroxygen disinfectant to minimise shoe sole contamination in a veterinary hospital. *Journal of the American Veterinary Medical Association* 228, 1391–1396.
- Barkema, H.W., Green, M.J., Bradley, A.J., Zadoks, R.N., 2009. The role of contagious disease in udder health. *Journal of Dairy Science* 92, 4717–4729.
- Barkema, H.W., Bartels, C., Wuickhuise, van L., Hesselink, J., Holzhauser, M., Weber, M., Franken, P., Kock, P., Brusckhe, C., Zimmer, G., 2001. Outbreak of bovine virus diarrhoea on Dutch dairy farms induced by a bovine herpesvirus 1 marker vaccine contaminated with bovine virus diarrhoea virus type 2. *Tijdschrift voor Diergeneeskunde* 126, 158–165.
- Barrett, D.J., Mee, J.F., Mullowney, P., Good, M., McGrath, G., Clegg, T., More, S.J., 2011. Risk factors associated with John's disease test status in dairy herds in Ireland. *Veterinary Record* 168, 410–411.
- Bazeley, K., 2009. Managing the risk of buying in disease when farmers buy in cattle. *UK Vet* 14, 42–45.
- Bazeley, K., 2011. Case report: *Campylobacter* infection, a beginner's mistakes. *UK Vet* 16, 29–30.
- Beard, P.M., Daniels, M.J., Henderson, D., Pirie, A., Rudge, K., Buxton, D., Rhind, S., Greig, A., Hutchings, M.R., McKendrick, I., Stevenson, K., Sharp, J.M., 2001. Paratuberculosis infection of nonruminant wildlife in Scotland. *Journal of Clinical Microbiology* 39, 1517–1521.
- Bergevoet, R.H.M., Van Schaik, G., Veling, J., Backus, G.B.C., Franken, P., 2009. Economic and epidemiological evaluation of *Salmonella* control in Dutch dairy herds. *Preventive Veterinary Medicine* 89, 1–7.
- Bishop, H., Erkelens, J., Van Winden, S., 2010. Indications of a relationship between buying-in policy and infectious diseases on dairy farms in Wales. *Veterinary Record* 167, 644–647.
- Björkman, C., Johansson, O., Stenlund, S., Holmdahl, O.J., Uggla, A., 1996. Neospora species infection in a herd of dairy cattle. *Journal of the American Veterinary Medical Association* 208, 1441–1444.
- Böhm, M., White, P.C.L., Chambers, J., Smith, L., Hutchins, M.R., 2007. Wild deer as a source of infection for livestock and humans in the UK. *The Veterinary Journal* 174, 260–276.
- Boelaert, F., Speybroeck, N., Kruif-de, A., Aerts, M., Burzykowski, T., Molenberghs, G., Berkvens, D.L., 2005. Risk factors for bovine herpesvirus-1 seropositivity. *Preventive Veterinary Medicine* 69, 285–295.
- Breidenbach, E., Hernandez, J., Engels, M., Stark, K., 2005. Risk assessment and cost/benefit analysis of a potential outbreak of IBR in cattle following a national show in Switzerland. In: *Proceedings of the 1st Swiss Buiatrics Congress, Bern, 19–21 October, 2005*, p. 55.
- Brennan, M., Kemp, R., Christley, R., 2008. Direct and indirect contacts between cattle farms in north-west England. *Preventive Veterinary Medicine* 84, 242–260.
- Bruwer, H., Bartels, C., van Schaik, G., 2010. Association of herd demographics and biosecurity with cattle longevity and udder health in dairy herds. In: *Proceedings of the XXVI World Buiatrics Congress, Santiago, Chile, p. 195*.
- Burnens, A.P., Bonnemain, P., Bruderer, U., Schalch, L., Audige, L., Grand, le D., Poumarat, F., Nicolet, J., 1999. Study on the seroprevalence of infections by *Mycoplasmata bovis* in Swiss cattle including epidemiological analysis of risk factors in a local population (Jura region). *Schweizer Archiv für Tierheilkunde* 141, 455–460.
- Byrne, W., Fagan, J., McCormack, M., 2001. *Mycoplasmata bovis* arthritis as a sequel to respiratory disease in bought-in weanling cattle in the Republic of Ireland. *Irish Veterinary Journal* 54, 516–519.
- Caldow, G., 2004. Biosecurity, does it have a place in the management of beef herds in the United Kingdom? *Cattle Practice* 12, 149–153.
- Caldow, G., 2009. Clinical forum – BVDV control and eradication. Part 2: Approaches to herd level control and national eradication. *UK Vet* 14, 20–27.
- Carpenter, T.E., Gardner, I.A., Collins, M.T., Whitlock, R.H., 2004. Effects of prevalence and testing by enzyme-linked immunosorbent assay and fecal culture on the risk of introduction of *Mycobacterium avium* subsp.

- paratuberculosis-infected cows into dairy herds. *Journal of Veterinary Diagnostic Investigation* 16, 31–38.
- Cernicchiaro, N., Pearl, D.L., Ghimire, S., Gyles, C.L., Johnson, R.P., LeJeune, J.T., Ziebell, K., McEwen, S.A., 2009. Risk factors associated with *Escherichia coli* O157:H7 in Ontario beef cow–calf operations. *Preventive Veterinary Medicine* 92, 106–115.
- Chi, J., Weersink, A., VanLeeuwen, J.A., Keefe, G.P., 2002a. The economics of controlling infectious diseases on dairy farms. *Canadian Journal of Agricultural Economics* 50, 237–256.
- Chi, J., VanLeeuwen, J.A., Weersink, A., Keefe, G.P., 2002b. Management factors related to seroprevalences to bovine viral diarrhoea virus, bovine-leukosis virus, *Mycobacterium avium* subspecies *paratuberculosis*, and *Neospora caninum* in dairy herds in the Canadian Maritimes. *Preventive Veterinary Medicine* 55, 57–68.
- Collins, M.T., Gardner, I.A., Garry, F.B., Roussel, A.J., Wells, S.J., 2006. Consensus recommendations on diagnostic testing for the detection of paratuberculosis in cattle in the United States. *Journal of the American Veterinary Medical Association* 229, 1912–1919.
- Cook, J.G., Holliman, A., 2004. Mastitis due to *Nocardia asteroides* in a UK dairy herd following restocking after FMD. *Veterinary Record* 154, 267–268.
- Crawshaw, M., Caldwell, G., Gunne, G., Rusbridge, S., 2002. Herd Biosecurity for Cattle. <http://www.sac.ac.uk/mainrep/pdfs/tn502herdbiosecurity.pdf> (accessed 5 July 2011).
- Daly, R., 2011. Biosecurity at the farm level: The role of extension in preventing animal disease introductions. *Journal of Dairy Science* 94, 672.
- Daszak, P., Cunningham, A.A., Hyatt, A.D., 2000. Emerging infectious diseases of wildlife – Threats to biodiversity and human health. *Science* 287, 443–449.
- Davison, H.C., Sayers, A.R., Smith, R.P., Pascoe, S.J.S., Davies, R.H., Weaver, J.P., Evans, S.J., 2006. Risk factors associated with the salmonella status of dairy farms in England and Wales. *Veterinary Record* 159, 871–880.
- Davison, H.C., Smith, R.P., Sayers, A.R., Evans, S.J., 2003. Dairy farm characteristics, including biosecurity, obtained during a cohort study in England and Wales. *Cattle Practice* 11, 299–310.
- DiGiacomo, R.F., Darlington, R.L., Evermann, J.F., 1985. Natural transmission of bovine leukemia virus in dairy calves by dehorning. *Canadian Journal of Comparative Medicine* 49, 340–342.
- Duncan, A.L., 1990. Health security in cattle herds. *In Practice* 12, 29–32.
- Dunowska, M., Morley, P.S., Patterson, G., Hyatt, D.R., Van Metre, D.C., 2006. Evaluation of the efficacy of a Peroxygen disinfectant-filled footmat for reduction of bacterial load on footwear in a large animal hospital setting. *Journal of the American Veterinary Medical Association* 228, 1935–1939.
- EFSA, 2006. Opinion of the Scientific Panel on Animal Health and Welfare (AHAW) on a request from the Commission on a Definition of a BoHV-1-free animal and a BoHV-1-free holding, and the procedures to verify and maintain this status. *EFSA Journal* 311, 1–65.
- Ellis-Iversen, J., Smith, R.P., Gibbins, J.C., Sharpe, C.E., Dominguez, M., Cook, A.J.C., 2011. Risk factors for transmission of foot-and-mouth disease during an outbreak in southern England in 2007. *Veterinary Record* 168, 128.
- Elvander, M., 1996. Severe respiratory disease in dairy cows caused by infection with bovine respiratory syncytial virus. *Veterinary Record* 138, 101–105.
- Evans, S.J., 1996. A case control study of multiple resistant *Salmonella* Typhimurium DT 104 infection of cattle in Great Britain. *Cattle Practice* 4, 259–266.
- Faust, M.A., Kinsel, M.L., Kirkpatrick, M.A., 2001. Characterizing biosecurity, health, and culling during dairy herd expansions. *Journal of Dairy Science* 84, 955–965.
- García-Bocanegra, I., Arenas-Montes, A., Lorca-Oro, C., Pujols, J., Gonzalez, M.A., Napp, S., Gómez-Guillamón, F., Zorrilla, I., Miguel, E.S., Arenas, A., 2011. Role of wild ruminants in the epidemiology of bluetongue virus serotypes 1, 4 and 8 in Spain. *Veterinary Research* 42, 88.
- Gilbert, M., Mitchell, A., Bourn, D., Mawdsley, J., Clifton-Hadley, R., Wint, W., 2005. Cattle movements and bovine tuberculosis in Great Britain. *Nature* 26, 491–496.
- Givens, M.D., Marley, S.D., 2008. Approaches to biosecurity in bovine embryo transfer programs. *Theriogenology* 69, 129–136.
- Godden, S., McMartin, S., Feirtag, J., Stabel, J., Bey, R., Goyal, S., Metzger, L., Fetrow, J., Wells, S., Chester-Jones, H., 2006. Heat-treatment of bovine colostrum. II: Effects of heating duration on pathogen viability and immunoglobulin G. *Journal of Dairy Science* 89, 3476–3483.
- Gopal, R., Goodchild, A., Hewinson, G., de la Rúa Domenech, R., Clifton-Hadley, R., 2006. Introduction of bovine tuberculosis to north-east England by bought-in cattle. *Veterinary Record* 159, 265–27.
- Greger, M., 2007. The long haul: Risks associated with livestock transport. *Biosecurity Bioterrorism* 5, 301–311.
- Greig, A., Stevenson, K., Henderson, D., Perez, V., Hughes, V., Pavlik, I., Hines II, M.E., McKendrick, I., Sharp, J.M., 1999. Epidemiological Study of Paratuberculosis in Wild Rabbits in Scotland. *Journal of Clinical Microbiology* 37, 1746–1751.
- Gunn, H.M., 1993. Role of fomites and flies in the transmission of bovine viral diarrhoea virus. *Veterinary Record* 132, 584–585.
- Gunn, G.J., Heffernan, C., Hall, M., McLeod, A., Hovi, M., 2008. Measuring and comparing constraints to improved biosecurity amongst GB farmers, veterinarians and the auxiliary industries. *Preventive Veterinary Medicine* 84, 310–323.
- Heffernan, C., Nielsen, L., Thomson, K., Gunn, G., 2008. An exploration of the drivers to bio-security collective action among a sample of UK cattle and sheep farmers. *Preventive Veterinary Medicine* 87, 358–372.
- Hemmingsen, B., Bitsch, V., 1983. Infection with IBR-virus (IBR/IPV-virus) in cattle. *Dansk Veterinaertidsskrift* 66, 939–943.
- Hilton, W.M., Olynk, N.J., 2011. Profitability of preconditioning: Lessons learned from an 11-year case study of an Indiana beef herd. *The Bovine Practitioner* 45, 40–50.
- Holliman, A., 2003. Restocking problems post-FMD. *Cattle Practice* 11, 311–312.
- Hopkins, S.G., DiGiacomo, R.F., 1997. Natural transmission of bovine leukemia virus in dairy and beef cattle. *Veterinary Clinics of North America Food Animal Practice* 13, 107–128.
- Jackson, R., Cooke, M.M., Coleman, J.D., Morris, R.S., 1995. Naturally occurring tuberculosis caused by *Mycobacterium bovis* in brushtail possums (*Trichosurus vulpecula*): I. An epidemiological analysis of lesion distribution. *New Zealand Veterinary Journal* 43, 306–314.
- Jones, T.O., 2011. Tail docking of dairy cattle in the UK. *Veterinary Record* 168, 542–543.
- King, J.S., Jenkins, D.J., Ellis, J.T., Fleming, P., Windsor, P.A., Lapeta, J., 2011. Implications of wild dog ecology on the sylvatic and domestic life cycle of *Neospora caninum* in Australia. *The Veterinary Journal* 188, 24–33.
- Kirk, J., Boggs, C., Jeffrey, J., Cardona, C., 2003. Efficacy of disinfectants for sanitizing boots under dairy farm conditions. *The Bovine Practitioner* 37, 50–53.
- Kristensen, E., Jakobsen, E.B., 2011. Danish farmers' perception of biosecurity. *Preventive Veterinary Medicine* 99, 122–129.
- Lang-Ree, J.R., Vatn, T., Kommisrud, E., Løken, T., 1994. Transmission of bovine viral diarrhoea virus by rectal examination. *Veterinary Record* 135, 412–413.
- Larson, E.L., 1995. APIC guideline for handwashing and hand antisepsis in health care settings. *American Journal of Infection and Control* 23, 251–269.
- Lindberg, A., Houe, H., 2005. Characteristics in the epidemiology of bovine viral diarrhoea virus (BVDV) of relevance to control. *Preventive Veterinary Medicine* 72, 55–73.
- Lindberg, A.L.E., Alenius, S., 1999. Principles for eradication of bovine viral diarrhoea virus (BVDV) infections in cattle populations. *Veterinary Microbiology* 64, 197–222.
- Majewska, A.M., Graczyk, T.K., Słodkiewicz-Kowalska, A., Tamang, L., Jędrzejewski, S., Zduniak, P., Solarczyk, P., Nowosad, A., Nowosad, P., 2009. The role of free-ranging, captive, and domestic birds of Western Poland in environmental contamination with *Cryptosporidium parvum* oocysts and *Giardia lamblia* cysts. *Parasitology Research* 104, 1093–1099.
- Makoschey, B., Beer, M., 2004. Assessment of the risk of transmission of vaccine viruses by using insufficiently cleaned injection devices. *Veterinary Record* 155, 563–564.
- Mathews, F., Lovett, L., Rushton, S., Macdonald, D.W., 2006. Bovine tuberculosis in cattle reduced risk on wildlife-friendly farms. *Biological Letters* 2, 271–274.
- Maunsell, F., Donovan, G.A., 2008. Biosecurity and risk management for dairy replacements. *Veterinary Clinics of North America Food Animal Practice* 24, 155–190.
- McLoughlin, M.F., McIlroy, S.G., Neill, S.D., 1998. A major outbreak of botulism in cattle being fed ensiled poultry litter. *Veterinary Record* 122, 579–581.
- Menzies, F.D., McCullough, S.J., McKeown, I.M., Forster, J.L., Jess, S., Batten, C., Murchie, A.K., Gloster, J., Fallows, J.G., Pelgrim, W., Mellor, P.S., Oura, C.A.L., 2008. Evidence for transplacental and contact transmission of bluetongue virus in cattle. *Veterinary Record* 163, 203–209.
- Moore, D.A., Kohrs, P., Baszler, T., Faux, C., Sathre, P., Wenz, J.R., Eldridge, L., Li, H., 2010. Outbreak of malignant catarrhal fever among cattle associated with a state livestock exhibition. *Journal of the American Veterinary Medical Association* 237, 87–92.
- Moore, D.A., Adaska, J.M., Higginbotham, G.E., Castillo, A.R., Collar, C.A., Sischo, W.M., 2009. Testing new dairy cattle for disease can boost herd health, cut costs. *California Agriculture* 63, 29–34.
- Moore, D.A., Merryman, M.L., Hartman, M.L., Klingborg, D.J., 2008. Comparison of published recommendations regarding biosecurity practices for various production animal species and classes. *Journal of the American Veterinary Medical Association* 233, 249–256.
- More, S.J., 2009. What is needed to eradicate bovine tuberculosis successfully: An Irish perspective. *The Veterinary Journal* 180, 275–278.
- More, S.J., Doherty, M.L., Downey, L., McKenzie, K., Devitt, C., O'Flaherty, J., 2011. Animal Health Ireland: Providing national leadership and coordination of non-regulatory animal health issues in Ireland. *Revue Scientifique et Technique Office International des Epizooties* 30, 715–723.
- Morley, P.S., 2002. Biosecurity of veterinary practices. *Veterinary Clinics of North America Food Animal Practice* 18, 133–155.
- Nickell, J.S., White, B.J., Larson, R.L., renter, D.G., Sanderson, M.W., Peck, C., 2011. Bovine viral diarrhoea virus (BVDV) status in cow–calf herds and associations with biosecurity and production practices among Montana beef producers. *Bovine Practitioner* 45, 14–22.
- Nicholson, F.A., Groves, S.J., Chambers, B.J., 2005. Pathogen survival during livestock manure storage and following land application. *Bioresource Technology* 96, 135–143.
- Nielsen, E.M., Tegtmeier, C., Andersen, H.J., Gronbaeck, C., Andersen, J.S., 2002. Influence of age, sex and herd characteristics on the occurrence of verocytotoxin-producing *Escherichia coli* O157 in Danish dairy farms. *Veterinary Microbiology* 88, 245–257.
- Niskanen, R., Lindberg, A., 2003. Transmission of bovine viral diarrhoea virus by unhygienic vaccination procedures, ambient air, and from contaminated pens. *The Veterinary Journal* 165, 125–130.
- Norton, S., Heuer, C., Jackson, R., 2009. A questionnaire-based cross-sectional study of clinical Johne's disease on dairy farms in New Zealand. *New Zealand Veterinary Journal* 57, 34–43.

- Nydam, D.V., Mohammed, H.O., 2005. Quantitative risk assessment of *Cryptosporidium* species infection in dairy calves. *Journal of Dairy Science* 88, 3932–3943.
- O'Farrell, K.J., Dillon, P., Mee, J.F., Crosse, S., Nolan, M., Byrne, N., Reidy, M., Flynn, F., Condon, T., 2001. Strategy for restocking of Moorepark after depopulation following bovine spongiform encephalopathy. *Irish Veterinary Journal* 54, 70–75.
- O'Corry Crowe, G., Hammond, R., Eves, J., Hayden, T.J., 1996. The effect of reduction in badger density on the spatial organisation and activity of badgers *Meles meles* L. in relation to farms in central Ireland. *Biology and Environment: Proceedings – Royal Irish Academy* 96b, 147–158.
- Ohlson, A., Heuer, C., Lockhart, C., Traven, M., Emanuelson, U., Alenius, S., 2010. Risk factors for seropositivity to bovine coronavirus and bovine respiratory syncytial virus in dairy herds. *Veterinary Record* 167, 201–207.
- Olea-Popelka, F.J., White, P.W., Collins, J.D., O'Keefe, J., Kelton, D.F., Martin, S.W., 2004. Breakdown severity during a bovine tuberculosis episode as a predictor of future herd breakdowns in Ireland. *Preventive Veterinary Medicine* 63, 163–172.
- Olde Riekerink, R.G.M., Barkema, H.W., Scholl, D.T., Poole, D.E., Kelton, D.F., 2010. Management practices associated with the bulk-milk prevalence of *Staphylococcus aureus* in Canadian dairy farms. *Preventive Veterinary Medicine* 97, 20–28.
- Oliveira, F.C.S., Azevedo, S.S., Pinheiro, S.R., Batista, C.S.A., Moraes, Z.M., Souza, G.O., Goncalves, A.P., Vasconcellos, S.A., 2010. Risk factors associated with leptospirosis in cows in the state of Bahia, northeastern Brazil. *Pesquisa Veterinaria Brasileira* 30, 398–402.
- Ortiz-Pelaez, A., Pfeiffer, D., 2008. Use of data mining techniques to investigate disease risk classification as a proxy for compromised biosecurity of cattle herds in Wales. *BMC Veterinary Research* 4, 24.
- Palmer, M.V., Whipple, D.L., Payeur, J.B., Alt, D.P., Esch, K.J., Bruning-Fann, C.S., Kaneene, J.B., 2000. Naturally occurring tuberculosis in white-tailed deer. *Journal of the American Veterinary Medical Association* 216, 1921–1924.
- Papini, R., Nardoni, S., Fanelli, A., Mancianti, F., 2009. High infection rate of *Trichophyton verrucosum* in calves from Central Italy. *Zoonoses and Public Health* 56, 59–64.
- Passchyn, P., Piepers, S., De Meulemeester, L., Boyen, F., Haesebrouck, F., De Vliegher, S., 2012. Between-herd prevalence of *Mycoplasma bovis* in bulk milk in Flanders, Belgium. *Research in Veterinary Science* 92, 219–220.
- Pillars, R.B., Grooms, D.L., Woltanski, J.A., Blair, E., 2009. Prevalence of Michigan dairy herds infected with *Mycobacterium avium* subspecies *paratuberculosis* as determined by environmental sampling. *Preventive Veterinary Medicine* 89, 191–196.
- Pritchard, G.C., 1996. Added animals: The challenge to preventive medicine. *Cattle Practice* 4, 253–257.
- Raaperi, K., Nurmoja, I., Orro, T., Viltrop, A., 2010. Seroepidemiology of bovine herpesvirus 1 (BHV1) infection among Estonian dairy herds and risk factors for the spread within herds. *Preventive Veterinary Medicine* 96, 74–81.
- Ramírez-Villaescusa, A.M., Medley, G.F., Mason, S., Green, L.E., 2010. Risk factors for herd breakdown with bovine tuberculosis in 148 cattle herds in the south west of England. *Preventive Veterinary Medicine* 95, 224–230.
- Richardson, E.K.B., Mee, J.F., Sanchez-Miguel, C., Crilly, J., More, S.J., 2009. Demographics of cattle positive for *Mycobacterium avium* subspecies *paratuberculosis* by faecal culture, from submissions to the Cork Regional Veterinary Laboratory. *Irish Veterinary Journal* 62, 398–405.
- Rodriguez-Lainz, A., Melendez-Retamal, P., Hird, D.W., Read, D.H., Walker, R.L., 1999. Farm- and host-level risk factors for papillomatous digital dermatitis in Chilean dairy cattle. *Preventive Veterinary Medicine* 42, 87–97.
- Samra, N.A., Jori, F., Samie, A., Thompson, P., 2011. The prevalence of *Cryptosporidium* spp. oocysts in wild mammals in the Kruger National Park, South Africa. *Veterinary Parasitology* 175, 155–159.
- Sanderson, M.W., Dargatz, D.G., Garry, F.B., 2000. Biosecurity practices of beef cow-calf producers. *Journal of the American Veterinary Medical Association* 217, 185–189.
- Santman-Berends, I.M.G.A., van Wuijckhuise, L., Vellema, P., van Rijn, P.A., 2010. Vertical transmission of bluetongue virus serotype 8 virus in Dutch dairy herds in 2007. *Veterinary Microbiology* 141, 31–35.
- Scharko, P.B., Giles, R.C., Vickers, M.L., McCoy, M.H., 2011. Diagnosis of congenital bovine viral diarrhoea infection in beef cow-calf herd from fence-line contact. *Proceedings of the American Association of Bovine Practitioners* 44, 147.
- Schouten, J.M., Bouwknegt, M., van de Giessen, A.W., Frankena, K., De Jong, M.C.M., Graat, E.A.M., 2004. Prevalence estimation and risk factors for *Escherichia coli* O157 on Dutch dairy farms. *Preventive Veterinary Medicine* 64, 49–61.
- Sibley, R., 2010. Biosecurity in the dairy herd. *In Practice* 32, 274–280.
- Simpson, V.R., 2002. Wild animals as reservoirs of infectious diseases in the UK. *The Veterinary Journal* 163, 128–146.
- Stevens, E.T., Thomson, D.U., Wileman, B.W., O'Dell, S., Chase, C.C.L., 2011. The survival of bovine viral diarrhoea virus on materials associated with livestock production. *The Bovine Practitioner* 45, 118–123.
- Sturdee, A.P., Chalmers, R.M., Bull, S.A., 1999. Detection of *Cryptosporidium* in wild animals in Britain. *Veterinary Parasitology* 80, 273–280.
- Sutmoller, P., Barteling, S.S., Olascoaga, R.C., Sumption, K.J., 2003. Control and eradication of foot-and-mouth disease. *Virus Research* 91, 101–144.
- Truyers, I.G.R., Mellor, D.J., Norquay, R., Gunn, G.J., Ellis, K.A., 2011. The eradication programme for bovine viral diarrhoea virus (BVDV) in Orkney 2001–2008. *Cattle Practice* 19, 47.
- Utenthal, A., Jensen, N.P.B., Blom, J.Y., 1996. Viral aetiology of enzootic pneumonia in Danish dairy herds: Diagnostic tools and epidemiology. *Veterinary Record* 139, 114–117.
- Valle, P., Martin, S.W., Tremblay, R., Bateman, K., 1999. Factors associated with being a bovine-virus diarrhoea (BVD) seropositive dairy herd in the Møre and Romsdal County of Norway. *Preventive Veterinary Medicine* 40, 165–177.
- Van Schaik, G., Schukken, Y.H., Nielen, M., Dijkhuizen, A.A., Benedictus, G., 2001. Risk factors for introduction of BHV1 into BHV1-free Dutch dairy farms: a case-control study. *Veterinary Quarterly* 23, 71–76.
- Van Schaik, G., Schukken, Y.H., Nielen, M., Dijkhuizen, A.A., Barkema, H.W., Benedictus, G., 2002. Probability of and risk factors for introduction of infectious diseases into Dutch SPF dairy farms: a cohort study. *Preventive Veterinary Medicine* 54, 279–289.
- Van Winden, S., Stevens, K., Guitian, J., McGowan, M., 2005. Preliminary findings of a systematic review and expert opinion workshop on biosecurity on cattle farms in the UK. *Cattle Practice* 13, 135–140.
- Van Wuijckhuise, L., Bosch, J., Franken, P., Frankena, K., Elbers, A.R.W., 1998. Epidemiological characteristics of bovine herpesvirus 1 infections determined by bulk milk testing of all Dutch dairy herds. *Veterinary Record* 142, 181–184.
- Velthuis, A.G., Mourits, M.C., 2007. Effectiveness of movement-prevention regulations to reduce the spread of foot-and-mouth disease in The Netherlands. *Preventive Veterinary Medicine* 82, 262–281.
- Ward, A.I., Judge, J., Delahay, R.J., 2010. Farm husbandry and badger behaviour: Opportunities to manage badger to cattle transmission of *Mycobacterium bovis*? *Preventive Veterinary Medicine* 93, 2–10.
- Weering, H.v., Mars, M., Muskens, J., Middeltesch, H., Schaik, G.v., 2005. The effect of biosecurity measures for paratuberculosis on the seroprevalence in Dutch dairy herds. In: *Proceedings of 8ICP 2005, Theme 3a: Prevention and Control – Herd Level*, pp. 246–253.
- Wells, S.J., 2000. Biosecurity on dairy operations: Hazards and risks. *Journal of Dairy Science* 83, 2380–2386.
- Whittington, R.J., Marshall, D.J., Nicholls, P.J., Marsh, I.B., Reddacliff, L.A., 2004. Survival and dormancy of *Mycobacterium avium* subsp. *paratuberculosis* in the environment. *Applied Environmental Microbiology* 70, 2989–3004.
- Williams, D.J.L., Guy, C.S., McGarry, J.W., Guy, F., Tasker, L., Smith, R.F., MacEachern, K., Cripps, P.J., Kelly, D.F., Trees, A.J., 2000. *Neospora caninum*-associated abortion in cattle: The time of experimentally-induced parasitaemia during gestation determines fetal survival. *Parasitology* 121, 347–358.
- Wolfe, D.M., Berke, O., Kelton, D.F., White, P.W., More, S.J., O'Keefe, J., Martin, S.W., 2010. From explanation to prediction: Developing a predictive model for recurrent bovine tuberculosis in Irish cattle herds. *Preventive Veterinary Medicine* 94, 170–177.
- Woodbine, K.A., Medley, G.F., Moore, S.J., Ramirez-Villaescusa, A., Mason, S., Green, L.E., 2008. A four year longitudinal sero-epidemiology study of *Neospora caninum* in adult cattle from 114 cattle herds in south west England: Associations with age, herd and dam-offspring pairs. *BMC Veterinary Research* 4. <http://dx.doi.org/10.1186/1746-6148-4-35>.